

ENERGY-SUPERCritical NLS: CRITICAL $\dot{H}^s$ -BOUNDS IMPLY SCATTERING

ROWAN KILLIP AND MONICA VISAN

ABSTRACT. We consider two classes of defocusing energy-supercritical nonlinear Schrödinger equations in dimensions $d \geq 5$. We prove that if the solution u is apriorily bounded in the critical Sobolev space, that is, $u \in L_t^\infty \dot{H}_x^{s_c}$, then u is global and scatters.

1. INTRODUCTION

We consider the initial-value problem for the defocusing nonlinear Schrödinger equation in dimension $d \geq 5$,

$$\begin{cases} iu_t = -\Delta u + F(u) \\ u(t=0, x) = u_0(x), \end{cases} \quad (1.1)$$

where the nonlinearity $F(u) = |u|^p u$ is energy-supercritical, that is, $p > \frac{4}{d-2}$.

The class of solutions to (1.1) is left invariant by the scaling

$$u(t, x) \mapsto \lambda^{\frac{2}{p}} u(\lambda^2 t, \lambda x). \quad (1.2)$$

This defines a notion of *criticality*. More precisely, a quick computation shows that the only homogeneous L_x^2 -based Sobolev space left invariant by the scaling is $\dot{H}_x^{s_c}(\mathbb{R}^d)$, where the *critical regularity* is $s_c := \frac{d}{2} - \frac{2}{p}$. If the regularity of the initial data to (1.1) is higher/lower than the critical regularity s_c , we call the problem *subcritical/supercritical*.

We consider (1.1) for initial data belonging to the critical homogeneous Sobolev space, that is, $u_0 \in \dot{H}_x^{s_c}(\mathbb{R}^d)$, in two regimes where $s_c > 1$ and $d \geq 5$. We prove that any maximal-lifespan solution that is uniformly bounded (throughout its lifespan) in $\dot{H}_x^{s_c}(\mathbb{R}^d)$ must be global and scatter. We were prompted to consider this problem by a recent preprint of Kenig and Merle [15] which proves similar results for radial solutions to the nonlinear wave equation in $\mathbb{R}^3$.

Let us start by making the notion of a solution more precise.

Definition 1.1 (Solution). A function $u : I \times \mathbb{R}^d \rightarrow \mathbb{C}$ on a non-empty time interval $0 \in I \subset \mathbb{R}$ is a *solution* (more precisely, a strong $\dot{H}_x^{s_c}(\mathbb{R}^d)$ solution) to (1.1) if it lies in the class $C_t^0 \dot{H}_x^{s_c}(K \times \mathbb{R}^d) \cap L_{t,x}^{p(d+2)/2}(K \times \mathbb{R}^d)$ for all compact $K \subset I$, and obeys the Duhamel formula

$$u(t) = e^{it\Delta} u(0) - i \int_0^t e^{i(t-s)\Delta} F(u(s)) ds \quad (1.3)$$

for all $t \in I$. We refer to the interval I as the *lifespan* of u . We say that u is a *maximal-lifespan solution* if the solution cannot be extended to any strictly larger interval. We say that u is a *global solution* if $I = \mathbb{R}$.

We define the *scattering size* of a solution to (1.1) on a time interval I by

$$S_I(u) := \int_I \int_{\mathbb{R}^d} |u(t, x)|^{\frac{p(d+2)}{2}} dx dt.$$

Associated to the notion of solution is a corresponding notion of blowup. As we will see in Theorem 1.6 below, this precisely corresponds to the impossibility of continuing the solution.

Definition 1.2 (Blowup). We say that a solution u to (1.1) *blows up forward in time* if there exists a time $t_1 \in I$ such that

$$S_{[t_1, \sup I)}(u) = \infty$$

and that u *blows up backward in time* if there exists a time $t_1 \in I$ such that

$$S_{(\inf I, t_1]}(u) = \infty.$$

We subscribe to the following conjecture.

Conjecture 1.3. *Let $d \geq 1$, $p \geq \frac{4}{d}$, and let $u : I \times \mathbb{R}^d \rightarrow \mathbb{C}$ be a maximal-lifespan solution to (1.1) such that $u \in L_t^\infty \dot{H}_x^{s_c}$. Then u is global and moreover,*

$$S_{\mathbb{R}}(u) \leq C(\|u\|_{L_t^\infty \dot{H}_x^{s_c}}), \quad (1.4)$$

for some function $C : [0, \infty) \rightarrow [0, \infty)$.

Our primary goal in this paper is to demonstrate how techniques developed to treat the energy-critical NLS can be applied to Conjecture 1.3 in the regime $s_c \geq 1$, although some of the arguments we will use were developed first in the mass-critical setting. As we will describe, the appearance of the $L_t^\infty \dot{H}_x^{s_c}$ norm on the right-hand side of (1.4) renders illusory the supercriticality of the equation. The famed supercriticality of Navier–Stokes is the fact that the problem is supercritical with respect to all quantities controlled by (known) conservation/monotonicity laws; see, for instance, the discussion in [31]. In the context of Conjecture 1.3, the assumption that the solution is uniformly bounded in $\dot{H}_x^{s_c}$ plays the role of the missing critical conservation law. It is not surprising therefore that techniques developed to treat problems with true critical conservation laws should be applicable in this setting. Next we review some of this work before describing the particular contribution of this paper.

Mass and energy are the only known coercive conserved quantities for NLS; hence, the corresponding critical NLS equations have received the most attention. In the mass-critical case, the critical regularity is $s_c = 0$ (i.e. $p = \frac{4}{d}$) and the scaling (1.2) leaves the mass invariant, that is, the conserved quantity

$$M(u) := \int_{\mathbb{R}^d} |u(t, x)|^2 dx.$$

Similarly, in the energy-critical case, the critical regularity is $s_c = 1$ (i.e. $p = \frac{4}{d-2}$, $d \geq 3$) and the scaling (1.2) leaves invariant the energy,

$$E(u) := \int_{\mathbb{R}^d} [|\nabla u(t, x)|^2 + \frac{1}{2+p}|u(t, x)|^{2+p}] dx,$$

which is also a conserved quantity for (1.1).

In the defocusing energy-critical case, it is known that all $\dot{H}_x^1$ initial data lead to global solutions with finite scattering size. Indeed, this was proved by Bourgain [2], Grillakis [11], and Tao [28] for spherically-symmetric initial data, and by Colliander–Keel–Staffilani–Takaoka–Tao [8], Ryckman–Visan [24], and Visan [37, 38] for arbitrary initial data. For results in the focusing case see [13, 19].

Unlike for the nonlinear wave equation (NLW), all known monotonicity formulae for NLS (that is, Morawetz-type inequalities) scale differently than the energy. Ultimately, the ingenious induction on energy technique of Bourgain (and the concomitant identification of bubbles) introduces a length scale to the problem which makes it possible to use

non-invariantly scaling monotonicity formulae. All subsequent work has built upon this foundational insight.

The Lin–Strauss Morawetz inequality used by Bourgain has $\dot{H}_x^{1/2}$ -scaling and is best adapted to the spherically-symmetric problem. To treat the non-radial problem, Colliander, Keel, Staffilani, Takaoka, and Tao introduced an interaction Morawetz inequality; this has $\dot{H}_x^{1/4}$ -scaling, which is even further from the $\dot{H}_x^1$ -scaling of the energy-critical problem. As the reader may notice, in both cases the regularity associated to the monotonicity formulae is lower than the critical regularity of the equation. This is the philosophical basis of our belief that the techniques developed for treating the energy-critical problem should be broadly applicable to Conjecture 1.3 whenever $s_c \geq 1/2$. In this paper, we will by no means complete this program, but rather have chosen to present some selected results that give the flavor of our main thesis without becoming swamped with technicalities.

We turn our attention now to the defocusing mass-critical NLS. In this case, Conjecture 1.3 has been proved for spherically-symmetric L_x^2 initial data in all dimensions $d \geq 2$; see [18, 21, 35]. For a proof of the corresponding conjecture in the focusing case (for spherically-symmetric L_x^2 data with mass less than that of the ground state and $d \geq 2$) see [18, 21]. At present, we do not know how to deal with the Galilean symmetry possessed by this equation, except through suppressing it by assuming spherical symmetry. We also note that in this case, one needs to prove additional regularity (rather than decay) to gain access to the known monotonicity formulae.

The first instance of Conjecture 1.3 to fall at non-conserved critical regularity was the case $s_c = 1/2$ in dimension $d = 3$. This was achieved by Kenig and Merle, [14]. They used the concentration-compactness technique in the manner they pioneered in [13] together with the Lin–Strauss Morawetz inequality.

The present paper is motivated by a recent preprint of Kenig and Merle, [15], who consider spherically-symmetric solutions to a class of defocusing energy-supercritical nonlinear wave equations in three dimensions. They prove that if the solution is known to be uniformly bounded in the critical $\dot{H}_x^s$ -space throughout its lifetime, then the solution must be global and it must scatter; this is precisely the NLW analogue of Conjecture 1.3.

Earlier, we drew a parallel to the Navier–Stokes equation. The most natural analogue of Conjecture 1.3 in that setting is to show that boundedness of a critical norm implies global regularity. Such results are known; see [9] and the references therein.

In this paper we prove Conjecture 1.3 in several instances of defocusing energy-supercritical nonlinear Schrödinger equations in dimensions $d \geq 5$ for arbitrary (not necessarily spherically-symmetric) initial data. The approach we take is modeled after [19], which considers the energy-critical problem in dimensions $d \geq 5$ in both the defocusing and focusing cases.

We will consider two different settings. In the first one, the nonlinearity is cubic, that is, $F(u) = |u|^2 u$, and hence the critical regularity is $s_c = \frac{d-2}{2}$. Note that in dimension $d \geq 5$, this problem is energy-supercritical.

The second problem we consider is that of general (not necessarily polynomial) energy-supercritical nonlinearities in dimension $d \geq 5$, that is, $p > \frac{4}{d-2}$. In this case, we also impose some additional constraints on the power p . First, we ask that the nonlinearity obeys a certain smoothness condition; more precisely, we ask that $s_c < 1 + p$, which is equivalent to $2p^2 - p(d-2) + 4 > 0$. The role of this constraint is to allow us to take s_c -many derivatives of the nonlinearity $F(u)$; this is important in the development of the local theory. Moreover, in Section 6, we require that s_c and p obey some further constraints. Together, these amount to

$$\begin{cases} 1 < s_c < \frac{3}{2} & \text{for } d = 5, 6 \\ 1 < s_c < \frac{d+2-\sqrt{(d-2)^2-16}}{4} & \text{for } d \geq 7. \end{cases} \quad (1.5)$$

One should not view (1.5) as a major constraint on the size of the critical regularity s_c . Indeed, we claim that an interpolation of the techniques we present to treat the two problems outlined above can be used to treat any defocusing energy-supercritical NLS (with the solution apriorily bounded in $\dot{H}_x^{s_c}$) in dimensions $d \geq 5$, without any additional constraint on s_c if the power p is an even integer and requiring merely the smoothness condition $s_c < 1 + p$ for arbitrary powers p . However, for the sake of readability, we chose not to work in this greater generality.

Our main results are the following:

Theorem 1.4 (Spacetime bounds – the cubic). *Let $d \geq 5$ and $F(u) = |u|^2 u$. Let $u : I \times \mathbb{R}^d \rightarrow \mathbb{C}$ be a maximal-lifespan solution to (1.1) such that $u \in L_t^\infty \dot{H}_x^{\frac{d-2}{2}}(I \times \mathbb{R}^d)$. Then u is global and moreover,*

$$S_{\mathbb{R}}(u) \leq C(\|u\|_{L_t^\infty \dot{H}_x^{\frac{d-2}{2}}}).$$

Theorem 1.5 (Spacetime bounds). *Let $d \geq 5$ and assume the critical regularity s_c satisfies (1.5). Let $u : I \times \mathbb{R}^d \rightarrow \mathbb{C}$ be a maximal-lifespan solution to (1.1) such that $u \in L_t^\infty \dot{H}_x^{s_c}(I \times \mathbb{R}^d)$. Then u is global and moreover,*

$$S_{\mathbb{R}}(u) \leq C(\|u\|_{L_t^\infty \dot{H}_x^{s_c}}).$$

As we already mentioned, the proofs of Theorems 1.4 and 1.5 follow closely the approach taken in [19] to study the energy-critical problem. We outline the argument in subsection 1.1 below. Our decision to work in dimensions $d \geq 5$ was motivated by the fact that it allows us to employ some techniques used in [19], in particular, the double Duhamel trick. The natural approach in lower dimensions would be to use the frequency localized interaction Morawetz inequality in the spirit of [8]. While subsequent developments (some of which are reviewed in this paper) lead to simplifications of [8], this would still be a significant undertaking and we do not pursue it here.

The arguments presented here apply mutis mutandis to the corresponding Hartree equations; indeed, the fact that the nonlinearity depends polynomially on u for that equation means that it resembles the simpler cubic case treated here. We also believe that the arguments adapt to the corresponding energy-supercritical wave equations in dimensions $d \geq 6$; however, we have not worked through the details.

To study the global theory for (1.1), we must first develop a local theory for this equation. To this end, we revisit arguments by Cazenave and Weissler, [4], who treated the case $0 \leq s_c \leq 1$, as well as more sophisticated stability results in the spirit of [23, 32].

Theorem 1.6 (Local well-posedness). *Let d and s_c be as in Theorem 1.4 or 1.5. Then, given $u_0 \in \dot{H}_x^{s_c}(\mathbb{R}^d)$ and $t_0 \in \mathbb{R}$, there exists a unique maximal-lifespan solution $u : I \times \mathbb{R}^d \rightarrow \mathbb{C}$ to (1.1) with initial data $u(t_0) = u_0$. This solution also has the following properties:*

- (Local existence) I is an open neighbourhood of t_0 .
- (Blowup criterion) If $\sup I$ is finite, then u blows up forward in time (in the sense of Definition 1.2). If $\inf I$ is finite, then u blows up backward in time.
- (Scattering) If $\sup I = +\infty$ and u does not blow up forward in time, then u scatters forward in time, that is, there exists a unique $u_+ \in \dot{H}_x^{s_c}(\mathbb{R}^d)$ such that

$$\lim_{t \rightarrow +\infty} \|u(t) - e^{it\Delta} u_+\|_{\dot{H}_x^{s_c}(\mathbb{R}^d)} = 0. \quad (1.6)$$

Conversely, given $u_+ \in \dot{H}_x^{s_c}(\mathbb{R}^d)$ there is a unique solution to (1.1) in a neighbourhood of infinity so that (1.6) holds.

- (Small data global existence) If $\| |\nabla|^{s_c} u_0 \|_2$ is sufficiently small (depending on d, p), then u is a global solution which does not blow up either forward or backward in time. Indeed, in this case $S_{\mathbb{R}}(u) \lesssim \| |\nabla|^{s_c} u_0 \|_2^{p(d+2)/2}$.

In Section 3 we establish this theorem as a corollary of our stability results Theorems 3.3 and 3.4. These stability results are essential to the arguments we present, more specifically, to the proof of Theorem 1.12.

1.1. Outline of the proofs of Theorems 1.4 and 1.5. We argue by contradiction. We show that if either Theorem 1.4 or Theorem 1.5 failed, this would imply the existence of a very special type of counterexample. Such counterexamples are then shown to have a wealth of properties not immediately apparent from their construction, so many properties, in fact, that they cannot exist.

While we will make some further reductions later, the main property of the special counterexamples is almost periodicity modulo symmetries:

Definition 1.7 (Almost periodicity modulo symmetries). Suppose $s_c > 0$. A solution u to (1.1) with lifespan I is said to be *almost periodic modulo symmetries* if there exist functions $N : I \rightarrow \mathbb{R}^+$, $x : I \rightarrow \mathbb{R}^d$, and $C : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that for all $t \in I$ and $\eta > 0$,

$$\int_{|x-x(t)| \geq C(\eta)/N(t)} |\nabla|^{s_c} u(t, x)|^2 dx + \int_{|\xi| \geq C(\eta)N(t)} |\xi|^{2s_c} |\hat{u}(t, \xi)|^2 d\xi \leq \eta.$$

We refer to the function N as the *frequency scale function* for the solution u , x the *spatial center function*, and to C as the *compactness modulus function*.

Remark 1.8. The parameter $N(t)$ measures the frequency scale of the solution at time t , while $1/N(t)$ measures the spatial scale. It is possible to multiply $N(t)$ by any function of t that is bounded both above and below, provided that we also modify the compactness modulus function C accordingly.

Remark 1.9. When $s_c = 0$ the equation admits a new symmetry, namely, Galilei invariance. This introduces a *frequency center function* $\xi(t)$ in the definition of almost periodicity modulo symmetries; see [17, 34] for further discussion.

Remark 1.10. By the Ascoli–Arzela Theorem, a family of functions is precompact in $\dot{H}_x^{s_c}(\mathbb{R}^d)$ if and only if it is norm-bounded and there exists a compactness modulus function C so that

$$\int_{|x| \geq C(\eta)} |\nabla|^{s_c} f(x)|^2 dx + \int_{|\xi| \geq C(\eta)} |\xi|^{2s_c} |\hat{f}(\xi)|^2 d\xi \leq \eta$$

for all functions f in the family. Thus, an equivalent formulation of Definition 1.7 is as follows: u is almost periodic modulo symmetries if and only if

$$\{u(t) : t \in I\} \subseteq \{\lambda^{\frac{2}{p}} f(\lambda(x + x_0)) : \lambda \in (0, \infty), x_0 \in \mathbb{R}^d, \text{ and } f \in K\}$$

for some compact subset K of $\dot{H}_x^{s_c}(\mathbb{R}^d)$.

Remark 1.11. A further consequence of compactness modulo symmetries is the existence of a function $c : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ so that

$$\int_{|x-x(t)| \leq c(\eta)/N(t)} |\nabla|^{s_c} u(t, x)|^2 dx + \int_{|\xi| \leq c(\eta)N(t)} |\xi|^{2s_c} |\hat{u}(t, \xi)|^2 d\xi \leq \eta$$

for all $t \in I$ and $\eta > 0$.

With these preliminaries out of the way, we can now describe the first major milestone in the proof of Theorems 1.4 and 1.5.

Theorem 1.12 (Reduction to almost periodic solutions). *Suppose that Theorem 1.4 (or Theorem 1.5) failed. Then there exists a maximal-lifespan solution $u : I \times \mathbb{R}^d \rightarrow \mathbb{C}$ to (1.1) such that $u \in L_t^\infty \dot{H}_x^{s_c}(I \times \mathbb{R}^d)$, u is almost periodic modulo symmetries, and u blows up both forward and backward in time. Moreover, u has minimal $L_t^\infty \dot{H}_x^{s_c}$ -norm among all blowup solutions, that is,*

$$\sup_{t \in I} \|\nabla|^{s_c} u(t)\|_2 \leq \sup_{t \in J} \|\nabla|^{s_c} v(t)\|_2$$

for all maximal-lifespan solutions $v : J \times \mathbb{R}^d \rightarrow \mathbb{C}$ that blow up in at least one time direction.

The reduction to almost periodic solutions was first realized by Keraani [17] in the context of the mass-critical NLS. This was adapted to the energy-critical case by Kenig and Merle [13].

Theorem 1.12 provides no information on the modulation parameters $x(t)$ and $N(t)$. In order to treat the mass-critical NLS in two dimensions, a further reduction was introduced in [18] where the behaviour of $N(t)$ is heavily constrained. This argument was adapted to the energy-critical case in [19]. This latter argument is directly applicable to the setting of this paper and yields

Theorem 1.13 (Three special scenarios for blowup). *Suppose that Theorem 1.4 (or Theorem 1.5) failed. Then there exists a maximal-lifespan solution $u : I \times \mathbb{R}^d \rightarrow \mathbb{C}$, which obeys $u \in L_t^\infty \dot{H}_x^{s_c}(I \times \mathbb{R}^d)$, is almost periodic modulo symmetries, and $S_I(u) = \infty$. Moreover, we can also ensure that the lifespan I and the frequency scale function $N : I \rightarrow \mathbb{R}^+$ match one of the following three scenarios:*

- I. (Finite-time blowup) We have that either $\sup I < \infty$ or $|\inf I| < \infty$.
- II. (Soliton-like solution) We have $I = \mathbb{R}$ and

$$N(t) = 1 \quad \text{for all } t \in \mathbb{R}.$$

- III. (Low-to-high frequency cascade) We have $I = \mathbb{R}$,

$$\inf_{t \in \mathbb{R}} N(t) \geq 1, \quad \text{and} \quad \limsup_{t \rightarrow +\infty} N(t) = \infty.$$

Therefore, in order to prove Theorems 1.4 and 1.5 it suffices to preclude the existence of solutions that satisfy the criteria in Theorem 1.13. Following [19], the key step in all three scenarios above is to prove negative regularity, that is, the solution u lies in L_x^2 or better. In scenarios II and III, the proof that $u \in L_x^2$ requires $d \geq 5$; note that this expresses better decay of the solution at spatial infinity. Similar in spirit to [18, 19, 21], negative regularity is deduced (via almost periodicity) from the minimality of the solution considered; recall that u has minimal $L_t^\infty \dot{H}_x^{s_c}$ norm among all blowup solutions.

A further manifestation of this minimality is the absence of a scattered wave at the endpoints of the lifespan I ; more formally, we have the following Duhamel formulae, which play an important role in proving negative regularity. For a proof, see [34, Section 6] or [20, Section 5].

Lemma 1.14 (No waste Duhamel formulae). *Let u be an almost periodic solution to (1.1) on its maximal-lifespan I . Then, for all $t \in I$,*

$$\begin{aligned} u(t) &= \lim_{T \nearrow \sup I} i \int_t^T e^{i(t-t')\Delta} F(u(t')) dt' \\ &= - \lim_{T \searrow \inf I} i \int_T^t e^{i(t-t')\Delta} F(u(t')) dt', \end{aligned} \tag{1.7}$$

as weak limits in $\dot{H}_x^{s_c}$.

We preclude the finite-time blowup scenario in Section 5. In order to achieve this, we use an argument based on the Strichartz inequality and we take advantage of the finiteness of the blowup time. We prove that in this case, the solution must have zero mass/energy. This contradicts the fact that the solution blows up.

To preclude the remaining two scenarios, we follow closely the strategy in [19]. As described above, the main point is to prove additional decay/negative regularity; this is achieved in two steps. First, we prove that the solution belongs to $L_t^\infty L_x^q$ for certain values of q less than $dp/2$, which is the exponent obtained by applying Sobolev embedding to $\dot{H}_x^{s_c}$. Just as in [19], the proof of this first step involves a bootstrap argument built off the

Duhamel formulae (1.7). In order to disentangle frequency interactions, we make use of an ‘acausal’ Gronwall inequality, Lemma 2.13. In the second step, we upgrade this breach of scaling to negative regularity in L_x^2 -based spaces. To do this, we take advantage of the global existence together with a double Duhamel trick in the spirit of [19, 29]. In order to make the associated time integrals converge, we need both $d \geq 5$ and the decay proved in step one.

In Section 7, we use the negative regularity proved in Section 6 together with the conservation of mass to preclude the low-to-high frequency cascade.

In Section 8, we use the interaction Morawetz inequality to preclude the soliton. In order to do this, we need the negative regularity proved in Section 6.

Acknowledgements. The first author was supported by NSF grant DMS-0701085.

2. NOTATIONS AND USEFUL LEMMAS

2.1. Some notation. We write $X \lesssim Y$ or $Y \gtrsim X$ whenever $X \leq CY$ for some constant $C > 0$. We use $O(Y)$ to denote any quantity X such that $|X| \lesssim Y$. We use the notation $X \sim Y$ to mean $X \lesssim Y \lesssim X$. The fact that these constants depend upon the dimension d or the power p will be suppressed. If C depends upon some additional parameters, we will indicate this with subscripts; for example, $X \lesssim_u Y$ denotes the assertion that $X \leq C_u Y$ for some C_u depending on u ; similarly for $X \sim_u Y$, $X = O_u(Y)$, etc. We denote by $X \pm$ any quantity of the form $X \pm \varepsilon$ for any $\varepsilon > 0$.

For any spacetime slab $I \times \mathbb{R}^d$, we use $L_t^q L_x^r(I \times \mathbb{R}^d)$ to denote the Banach space of functions $u : I \times \mathbb{R}^d \rightarrow \mathbb{C}$ whose norm is

$$\|u\|_{L_t^q L_x^r(I \times \mathbb{R}^d)} := \left(\int_I \|u(t)\|_{L_x^r}^q dt \right)^{\frac{1}{q}} < \infty,$$

with the usual modifications when q or r is equal to infinity. When $q = r$ we abbreviate $L_t^q L_x^q$ as $L_{t,x}^q$.

We define the Fourier transform on $\mathbb{R}^d$ by

$$\hat{f}(\xi) := (2\pi)^{-d/2} \int_{\mathbb{R}^d} e^{-ix\xi} f(x) dx.$$

For $s \in \mathbb{R}$, we define the fractional differentiation/integral operator

$$|\nabla|^s \hat{f}(\xi) := |\xi|^s \hat{f}(\xi),$$

which in turn defines the homogeneous Sobolev norm

$$\|f\|_{\dot{H}_x^s(\mathbb{R}^d)} := \| |\nabla|^s f \|_{L_x^2(\mathbb{R}^d)}.$$

Finally, we use the notation

$$\nabla F(u(x)) := \nabla u(x) \cdot F'(u(x)) := \nabla u(x) F_z(u(x)) + \overline{\nabla u(x)} F_{\bar{z}}(u(x)),$$

where $F_z, F_{\bar{z}}$ denote the usual complex derivatives

$$F_z := \frac{1}{2} \left(\frac{\partial F}{\partial x} - i \frac{\partial F}{\partial y} \right), \quad F_{\bar{z}} := \frac{1}{2} \left(\frac{\partial F}{\partial x} + i \frac{\partial F}{\partial y} \right).$$

2.2. Strichartz estimates. Let $e^{it\Delta}$ be the free Schrödinger evolution. From the explicit formula

$$e^{it\Delta} f(x) = \frac{1}{(4\pi it)^{d/2}} \int_{\mathbb{R}^d} e^{i|x-y|^2/4t} f(y) dy,$$

one easily obtains the standard dispersive inequality

$$\|e^{it\Delta} f\|_{L_x^\infty(\mathbb{R}^d)} \lesssim |t|^{-\frac{d}{2}} \|f\|_{L_x^1(\mathbb{R}^d)} \quad (2.1)$$

for all $t \neq 0$. In particular, as the free propagator conserves the L_x^2 -norm,

$$\|e^{it\Delta}f\|_{L_x^p(\mathbb{R}^d)} \lesssim |t|^{d(\frac{1}{p}-\frac{1}{2})} \|f\|_{L_x^{p'}(\mathbb{R}^d)} \quad (2.2)$$

for all $t \neq 0$ and $2 \leq p \leq \infty$, where $\frac{1}{p} + \frac{1}{p'} = 1$.

Definition 2.1 (Admissible pairs). For $d \geq 3$, we say that a pair of exponents (q, r) is *Schrödinger-admissible* if

$$\frac{2}{q} + \frac{d}{r} = \frac{d}{2} \quad \text{and} \quad 2 \leq q, r \leq \infty. \quad (2.3)$$

For a fixed spacetime slab $I \times \mathbb{R}^d$, we define the *Strichartz norm*

$$\|u\|_{S^0(I)} := \sup_{(q,r) \text{ admissible}} \|u\|_{L_t^q L_x^r(I \times \mathbb{R}^d)}.$$

We write $S^0(I)$ for the closure of all test functions under this norm and denote by $N^0(I)$ the dual of $S^0(I)$.

A simple application of Sobolev embedding yields

$$\begin{aligned} \||\nabla|^{s_c} u\|_{L_t^\infty L_x^2(I \times \mathbb{R}^d)} + \||\nabla|^{s_c} u\|_{L_{t,x}^{\frac{2(d+2)}{d}}(I \times \mathbb{R}^d)} + \|u\|_{L_t^\infty L_x^{\frac{dp}{2}}(I \times \mathbb{R}^d)} + \|u\|_{L_{t,x}^{\frac{p(d+2)}{2}}(I \times \mathbb{R}^d)} \\ \lesssim \||\nabla|^{s_c} u\|_{S^0(I)} \end{aligned}$$

for all $d \geq 3$.

As a consequence of the dispersive estimate (2.1), we have the following standard Strichartz estimate.

Lemma 2.2 (Strichartz). *Let $s \geq 0$, let I be a compact time interval, and let $u : I \times \mathbb{R}^d \rightarrow \mathbb{C}$ be a solution to the forced Schrödinger equation*

$$iu_t + \Delta u = F.$$

Then,

$$\||\nabla|^s u\|_{S^0(I)} \lesssim \||\nabla|^s u(t_0)\|_{L_x^2} + \||\nabla|^s F\|_{N^0(I)}$$

for any $t_0 \in I$.

Proof. See, for example, [10, 27]. For the endpoint $(q, r) = (2, \frac{2d}{d-2})$ in dimensions $d \geq 3$, see [12]. $\square$

2.3. Basic harmonic analysis. Let $\varphi(\xi)$ be a radial bump function supported in the ball $\{\xi \in \mathbb{R}^d : |\xi| \leq \frac{11}{10}\}$ and equal to 1 on the ball $\{\xi \in \mathbb{R}^d : |\xi| \leq 1\}$. For each number $N > 0$, we define the Fourier multipliers

$$\begin{aligned} \widehat{P_{\leq N} f}(\xi) &:= \widehat{f_{\leq N}}(\xi) := \varphi(\xi/N) \hat{f}(\xi) \\ \widehat{P_{> N} f}(\xi) &:= \widehat{f_{> N}}(\xi) := (1 - \varphi(\xi/N)) \hat{f}(\xi) \\ \widehat{P_N f}(\xi) &:= \widehat{f_N}(\xi) := (\varphi(\xi/N) - \varphi(2\xi/N)) \hat{f}(\xi) \end{aligned}$$

and similarly $P_{< N}$ and $P_{\geq N}$. We also define

$$P_{M < \cdot \leq N} := P_{\leq N} - P_{\leq M} = \sum_{M < N' \leq N} P_{N'}$$

whenever $M < N$. We will usually use these multipliers when M and N are *dyadic numbers* (that is, of the form 2^n for some integer n); in particular, all summations over N or M are understood to be over dyadic numbers. Nevertheless, it will occasionally be convenient to allow M and N to not be a power of 2.

Like all Fourier multipliers, the Littlewood-Paley operators commute with the propagator $e^{it\Delta}$, as well as with differential operators such as $i\partial_t + \Delta$. We will use basic properties of these operators many many times, including

Lemma 2.3 (Bernstein estimates). *For $1 \leq r \leq q \leq \infty$,*

$$\begin{aligned} \|\nabla^{\pm s} P_N f\|_{L_x^r(\mathbb{R}^d)} &\sim N^{\pm s} \|P_N f\|_{L_x^r(\mathbb{R}^d)}, \\ \|P_{\leq N} f\|_{L_x^q(\mathbb{R}^d)} &\lesssim N^{\frac{d}{r} - \frac{d}{q}} \|P_{\leq N} f\|_{L_x^r(\mathbb{R}^d)}, \\ \|P_N f\|_{L_x^q(\mathbb{R}^d)} &\lesssim N^{\frac{d}{r} - \frac{d}{q}} \|P_N f\|_{L_x^r(\mathbb{R}^d)}. \end{aligned}$$

Lemma 2.4 (Product rule, [6]). *Let $s \in (0, 1]$ and $1 < r, r_1, r_2, q_1, q_2 < \infty$ such that $\frac{1}{r} = \frac{1}{r_1} + \frac{1}{r_2}$ for $i = 1, 2$. Then,*

$$\|\nabla^s(fg)\|_r \lesssim \|f\|_{r_1} \|\nabla^s g\|_{q_1} + \|\nabla^s f\|_{r_2} \|g\|_{q_2}.$$

We will also need the following fractional chain rule from [6]. For a textbook treatment, see [36, §2.4].

Lemma 2.5 (Fractional chain rule, [6]). *Suppose $G \in C^1(\mathbb{C})$, $s \in (0, 1]$, and $1 < q, q_1, q_2 < \infty$ are such that $\frac{1}{q} = \frac{1}{q_1} + \frac{1}{q_2}$. Then,*

$$\|\nabla^s G(u)\|_q \lesssim \|G'(u)\|_{q_1} \|\nabla^s u\|_{q_2}.$$

When the function G is no longer C^1 , but merely Hölder continuous, we have the following chain rule:

Lemma 2.6 (Fractional chain rule for a Hölder continuous function, [38]). *Let G be a Hölder continuous function of order $0 < p < 1$. Then, for every $0 < s < p$, $1 < q < \infty$, and $\frac{s}{p} < \sigma < 1$ we have*

$$\|\nabla^s G(u)\|_q \lesssim \|u\|^{p-\frac{s}{\sigma}}_{q_1} \|\nabla^\sigma u\|_{\frac{s}{\sigma} q_2}^{\frac{s}{\sigma}}, \quad (2.4)$$

provided $\frac{1}{q} = \frac{1}{q_1} + \frac{1}{q_2}$ and $(1 - \frac{s}{p\sigma})q_1 > 1$.

As a direct consequence of the two fractional chain rules above and interpolation, we have the following

Corollary 2.7. *Let $F(u) = |u|^p u$ and let $s \geq 0$ if p is an even integer or $0 \leq s < 1 + p$ otherwise. Then, on any spacetime slab $I \times \mathbb{R}^d$ we have*

$$\|\nabla^s F(u)\|_{N^0(I)} \lesssim \|\nabla^s u\|_{S^0(I)} \|u\|_{L_{t,x}^{\frac{p(d+2)}{2}}}^p$$

and

$$\|\nabla^s F(u)\|_{L_t^\infty L_x^{\frac{2d}{d+4}}} \lesssim \|\nabla^s u\|_{L_t^\infty L_x^2} \|u\|_{L_t^\infty L_x^{dp/2}}^p.$$

Revisiting the proof of Lemma 2.6, we obtain the following lemma. In actually, the result can be deduced directly from (2.4) and Lemma 2.4; however, the result (2.6) appearing in the proof will be needed in Section 4.

Lemma 2.8. *Let G be a Hölder continuous function of order $0 < p \leq 1$ and let $0 < s < \sigma p < p$. For $1 < q, q_1, q_2, r_1, r_2, r_3 < \infty$ such that $\frac{1}{q} = \frac{1}{q_1} + \frac{1}{q_2} = \frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3}$ we have*

$$\begin{aligned} &\|\nabla^s[w \cdot (G(u+v) - G(u))]\|_q \\ &\lesssim \|\nabla^s w\|_{q_1} \|v\|_{p q_2}^p + \|w\|_{r_1} \|v\|_{(p-\frac{s}{\sigma})r_2}^{p-\frac{s}{\sigma}} (\|\nabla^\sigma v\|_{\frac{s}{\sigma} r_3} + \|\nabla^\sigma u\|_{\frac{s}{\sigma} r_3})^{\frac{s}{\sigma}}, \end{aligned}$$

provided $(1-p)r_1, (p-\frac{s}{\sigma})r_2 > 1$.

Proof. In [26], Strichartz proved that for all Schwartz functions f , $1 < q < \infty$, and $0 < s < 1$,

$$\|\nabla^s f\|_{L_x^q} \sim \|\mathcal{D}_s(f)\|_{L_x^q},$$

where

$$\mathcal{D}_s(f)(x) := \left(\int_0^\infty \left| \int_{|y|<1} |f(x+ry) - f(x)| dy \right|^2 \frac{dr}{r^{1+2s}} \right)^{1/2}. \quad (2.5)$$

In view of this, the claim will follow from the pointwise inequality

$$\begin{aligned} \mathcal{D}_s(w \cdot [G(u+v) - G(u)]) \\ \lesssim \mathcal{D}_s(w)|v|^p + [M(|w|^{\frac{1}{1-p}})]^{1-p} [M(|v|)]^{p-\frac{p}{\sigma}} [\mathcal{D}_\sigma(u+v) + \mathcal{D}_\sigma(u)]^{\frac{p}{\sigma}}, \end{aligned} \quad (2.6)$$

where M denotes the Hardy–Littlewood maximal function.

As G is Hölder continuous of order p ,

$$\begin{aligned} & |(w \cdot [G(u+v) - G(u)])(x+ry) - (w \cdot [G(u+v) - G(u)])(x)| \\ & \leq |w(x+ry) - w(x)| |G(u+v)(x) - G(u)(x)| \\ & \quad + |w(x+ry)| |[G(u+v) - G(u)](x+ry) - [G(u+v) - G(u)](x)| \\ & \lesssim |w(x+ry) - w(x)| |v(x)|^p + H(x) \end{aligned} \quad (2.7)$$

with

$$H := |w(x+ry)| |[G(u+v) - G(u)](x+ry) - [G(u+v) - G(u)](x)|. \quad (2.8)$$

Note that the first term on the right-hand side of (2.7) gives rise to the term $\mathcal{D}_s(w)|v|^p$ in (2.6). Hence it remains to estimate the contribution of H . In order to achieve this, we estimate H in two different ways:

$$H \lesssim |w(x+ry)| \left[|v(x+ry)|^p + |v(x)|^p \right] \quad (2.9)$$

and also

$$H \lesssim |w(x+ry)| \left[|(u+v)(x+ry) - (u+v)(x)|^p + |u(x+ry) - u(x)|^p \right]. \quad (2.10)$$

Using Hölder's inequality and (2.9), we see that

$$\begin{aligned} \int_{A(x)}^\infty \left| \int_{|y|<1} H dy \right|^2 \frac{dr}{r^{1+2s}} & \lesssim [M(|w|^{\frac{1}{1-p}})(x)]^{2(1-p)} [M(|v|)(x)]^{2p} \int_{A(x)}^\infty \frac{dr}{r^{1+2s}} \\ & \lesssim [A(x)]^{-2s} [M(|w|^{\frac{1}{1-p}})(x)]^{2(1-p)} [M(|v|)(x)]^{2p}. \end{aligned}$$

The precise value of $A(x)$ will be determined below.

We now turn our attention to small values of r . Using (2.10) together with Hölder's inequality, we see that

$$\begin{aligned} \int_{|y|<1} H dy & \lesssim [M(|w|^{\frac{1}{1-p}})(x)]^{1-p} \left[\int_{|y|<1} |(u+v)(x+ry) - (u+v)(x)| \right. \\ & \quad \left. + |u(x+ry) - u(x)| dy \right]^p \end{aligned}$$

and so, applying Hölder's inequality again, we find

$$\begin{aligned} \int_0^{A(x)} \left| \int_{|y|<1} H dy \right|^2 \frac{dr}{r^{1+2s}} \\ \lesssim [M(|w|^{\frac{1}{1-p}})(x)]^{2(1-p)} [A(x)]^{2(\sigma p - s)} [\mathcal{D}_\sigma(u+v)(x) + \mathcal{D}_\sigma(u)(x)]^{2p}. \end{aligned}$$

Putting things together and optimizing the choice of $A(x)$, we derive (2.6). This finishes the proof of the lemma. $\square$

The next result is formally similar to Lemma 2.6. The proof is simple; see the appendix in [20]. It is used in the proof of Lemma 6.2.

Lemma 2.9 (Nonlinear Bernstein, [20]). *Let $G : \mathbb{C} \rightarrow \mathbb{C}$ be Hölder continuous of order $0 < p \leq 1$. Then*

$$\|P_N G(u)\|_{L_x^{q/p}(\mathbb{R}^d)} \lesssim N^{-p} \|\nabla u\|_{L_x^q(\mathbb{R}^d)}^p$$

for any $1 \leq q < \infty$.

2.4. Concentration compactness. In this subsection we record the linear profile decomposition statement which will lead to the reduction in Theorem 1.12. We first recall the symmetries of the equation (1.1) which fix the initial surface $t = 0$.

Definition 2.10 (Symmetry group). For any phase $\theta \in \mathbb{R}/2\pi\mathbb{Z}$, position $x_0 \in \mathbb{R}^d$, and scaling parameter $\lambda > 0$, we define a unitary transformation $g_{\theta, x_0, \lambda} : \dot{H}_x^{s_c}(\mathbb{R}^d) \rightarrow \dot{H}_x^{s_c}(\mathbb{R}^d)$ by

$$[g_{\theta, x_0, \lambda} f](x) := \lambda^{-\frac{2}{p}} e^{i\theta} f(\lambda^{-1}(x - x_0)).$$

Recall that $s_c := \frac{d}{2} - \frac{2}{p}$. Let G denote the collection of such transformations. For a function $u : I \times \mathbb{R}^d \rightarrow \mathbb{C}$, we define $T_{g_{\theta, x_0, \lambda}} u : \lambda^2 I \times \mathbb{R}^d \rightarrow \mathbb{C}$ where $\lambda^2 I := \{\lambda^2 t : t \in I\}$ by the formula

$$[T_{g_{\theta, x_0, \lambda}} u](t, x) := \lambda^{-\frac{2}{p}} e^{i\theta} u(\lambda^{-2} t, \lambda^{-1}(x - x_0)).$$

Note that if u is a solution to (1.1), then $T_g u$ is a solution to (1.1) with initial data gu_0 .

Remark 2.11. It is easy to verify that G is a group and that the map $g \mapsto T_g$ is a homomorphism. The map $u \mapsto T_g u$ maps solutions to (1.1) to solutions with the same scattering size as u , that is, $S(T_g u) = S(u)$. Furthermore, u is a maximal-lifespan solution if and only if $T_g u$ is a maximal-lifespan solution.

We are now ready to state the linear profile decomposition; in the generality needed here, this was proved in [25]. For $s_c = 0$ the linear profile decomposition was proved in [1, 3, 22], while for $s_c = 1$ it was established in [16].

Lemma 2.12 (Linear profile decomposition, [25]). *Fix $s_c > 0$ and let $\{u_n\}_{n \geq 1}$ be a sequence of functions bounded in $\dot{H}_x^{s_c}(\mathbb{R}^d)$. Then, after passing to a subsequence if necessary, there exist functions $\{\phi^j\}_{j \geq 1} \subset \dot{H}_x^{s_c}(\mathbb{R}^d)$, group elements $g_n^j \in G$, and times $t_n^j \in \mathbb{R}$ such that for all $J \geq 1$ we have the decomposition*

$$u_n = \sum_{j=1}^J g_n^j e^{it_n^j \Delta} \phi^j + w_n^J$$

with the following properties:

- $w_n^J \in \dot{H}_x^{s_c}(\mathbb{R}^d)$ and obey

$$\lim_{J \rightarrow \infty} \limsup_{n \rightarrow \infty} \|e^{it \Delta} w_n^J\|_{L_{t,x}^{\frac{p(d+2)}{2}}(\mathbb{R} \times \mathbb{R}^d)} = 0.$$

- For any $j \neq j'$,

$$\frac{\lambda_n^j}{\lambda_n^{j'}} + \frac{\lambda_n^{j'}}{\lambda_n^j} + \frac{|x_n^j - x_n^{j'}|^2}{\lambda_n^j \lambda_n^{j'}} + \frac{|t_n^j (\lambda_n^j)^2 - t_n^{j'} (\lambda_n^{j'})^2|}{\lambda_n^j \lambda_n^{j'}} \rightarrow \infty \quad \text{as } n \rightarrow \infty.$$

- For any $J \geq 1$ we have the decoupling properties:

$$\lim_{n \rightarrow \infty} \left[\|\nabla |^{s_c} u_n\|_2^2 - \sum_{j=1}^J \|\nabla |^{s_c} \phi^j\|_2^2 - \|\nabla |^{s_c} w_n^J\|_2^2 \right] = 0$$

and for any $1 \leq j \leq J$,

$$e^{-it_n^j \Delta} [(g_n^j)^{-1} w_n^J] \rightarrow 0 \quad \text{weakly in } \dot{H}_x^{s_c} \text{ as } n \rightarrow \infty.$$

2.5. A Gronwall inequality. Our last technical tool is a form of Gronwall's inequality that involves both the past and the future, 'acausal' in the terminology of [30]. We import it from [19], where it was used for precisely the same purpose as it will be here.

Lemma 2.13 (Acausal Gronwall inequality, [19]). *Given $\gamma > 0$, $0 < \eta < \frac{1}{2}(1 - 2^{-\gamma})$, and $\{b_k\} \in \ell^\infty(\mathbb{Z}^+)$, let $x_k \in \ell^\infty(\mathbb{Z}^+)$ be a non-negative sequence obeying*

$$x_k \leq b_k + \eta \sum_{l=0}^{\infty} 2^{-\gamma|k-l|} x_l \quad \text{for all } k \geq 0.$$

Then

$$x_k \lesssim \sum_{l=0}^k r^{|k-l|} b_l \quad \text{for all } k \geq 0$$

for some $r = r(\eta) \in (2^{-\gamma}, 1)$. Moreover, $r \downarrow 2^{-\gamma}$ as $\eta \downarrow 0$.

3. LOCAL WELL-POSEDNESS

In this section we develop the local well-posedness theory for (1.1). The arguments we use are inspired by previous work on nonlinear Schrödinger equations at critical regularity. For $s_c \in [0, 1]$ the standard local well-posedness theory (see Theorem 3.1 below) was established by Cazenave and Weissler, [4]; see also [5, 30]. For stability results (see Theorems 3.3 and 3.4 below) in the mass- and energy-critical settings (i.e. $s_c = 0, 1$), see [8, 24, 32, 34]. In this section, we mainly follow the exposition in [20], which revisits the local theory for the mass- and energy-critical NLS.

We start with the following standard local well-posedness result, for which one assumes that the initial data lies in the inhomogeneous critical Sobolev space. This assumption simplifies the proof and can be removed a posteriori by using the stability results proved below.

Theorem 3.1 (Standard local well-posedness). *Let $d \geq 1$, $s_c \geq 0$, and let $u_0 \in H_x^{s_c}(\mathbb{R}^d)$. Assume in addition that $s_c < 1 + p$ if p is not an even integer. Then there exists $\eta_0 = \eta_0(d) > 0$ such that if $0 < \eta \leq \eta_0$ and I is a compact interval containing zero such that*

$$\left\| |\nabla|^{s_c} e^{it\Delta} u_0 \right\|_{L_t^{p+2} L_x^{\frac{2d(p+2)}{2(d-2)+dp}}(I \times \mathbb{R}^d)} \leq \eta, \quad (3.1)$$

then there exists a unique solution u to (1.1) on $I \times \mathbb{R}^d$. Moreover, we have the bounds

$$\left\| |\nabla|^{s_c} u \right\|_{L_t^{p+2} L_x^{\frac{2d(p+2)}{2(d-2)+dp}}(I \times \mathbb{R}^d)} \leq 2\eta \quad (3.2)$$

$$\left\| |\nabla|^{s_c} u \right\|_{S^0(I \times \mathbb{R}^d)} \lesssim \left\| |\nabla|^{s_c} u_0 \right\|_{L_x^2} + \eta^{1+p} \quad (3.3)$$

$$\|u\|_{S^0(I \times \mathbb{R}^d)} \lesssim \|u_0\|_{L_x^2}. \quad (3.4)$$

Remark 3.2. By the Strichartz inequality, we know that

$$\left\| |\nabla|^{s_c} e^{it\Delta} u_0 \right\|_{L_t^{p+2} L_x^{\frac{2d(p+2)}{2(d-2)+dp}}(\mathbb{R} \times \mathbb{R}^d)} \lesssim \left\| |\nabla|^{s_c} u_0 \right\|_{L_x^2}.$$

Thus, (3.1) holds with $I = \mathbb{R}$ for initial data with sufficiently small norm. Alternatively, by the monotone convergence theorem, (3.1) holds provided I is chosen sufficiently small. Note that by scaling, the length of the interval I depends on the fine properties of u_0 , not only on its norm.

Proof. We will essentially repeat the standard argument from [4]; the fractional chain rule Lemma 2.5 leads to some simplifications.

The theorem follows from a contraction mapping argument. More precisely, using the Strichartz estimates from Lemma 2.2, we will show that the map $u \mapsto \Phi(u)$ defined by

$$\Phi(u)(t) := e^{it\Delta}u_0 - i \int_0^t e^{i(t-s)\Delta} F(u(s)) ds,$$

is a contraction on the set $B_1 \cap B_2$ where

$$\begin{aligned} B_1 &:= \left\{ u \in L_t^\infty H_x^{s_c}(I \times \mathbb{R}^d) : \|u\|_{L_t^\infty H_x^{s_c}(I \times \mathbb{R}^d)} \leq 2\|u_0\|_{H_x^{s_c}} + C(d, p)(2\eta)^{1+p} \right\} \\ B_2 &:= \left\{ u \in L_t^{p+2} W_x^{s_c, \frac{2d(p+2)}{2(d-2)+dp}}(I \times \mathbb{R}^d) : \|\nabla|^{s_c} u\|_{L_t^{p+2} L_x^{\frac{2d(p+2)}{2(d-2)+dp}}(I \times \mathbb{R}^d)} \leq 2\eta \right. \\ &\quad \left. \text{and } \|u\|_{L_t^{p+2} L_x^{\frac{2d(p+2)}{2(d-2)+dp}}(I \times \mathbb{R}^d)} \leq 2C(d, p)\|u_0\|_{L_x^2} \right\} \end{aligned}$$

under the metric given by

$$d(u, v) := \|u - v\|_{L_t^{p+2} L_x^{\frac{2d(p+2)}{2(d-2)+dp}}(I \times \mathbb{R}^d)}.$$

Here $C(d, p)$ denotes a constant that changes from line to line. Note that the norm appearing in the metric scales like L_x^2 . Note also that both B_1 and B_2 are closed (and hence complete) in this metric.

Using the Strichartz inequality followed by Corollary 2.7 and Sobolev embedding, we find that for $u \in B_1 \cap B_2$,

$$\begin{aligned} \|\Phi(u)\|_{L_t^\infty H_x^{s_c}(I \times \mathbb{R}^d)} &\leq \|u_0\|_{H_x^{s_c}} + C(d, p) \|\nabla|^{s_c} F(u)\|_{L_t^{\frac{p+2}{p+1}} L_x^{\frac{2d(p+2)}{2(d+2)+dp}}(I \times \mathbb{R}^d)} \\ &\leq \|u_0\|_{H_x^{s_c}} + C(d, p) \|\nabla|^{s_c} u\|_{L_t^{p+2} L_x^{\frac{2d(p+2)}{2(d-2)+dp}}(I \times \mathbb{R}^d)} \|u\|_{L_t^{p+2} L_x^{\frac{dp(p+2)}{4}}(I \times \mathbb{R}^d)}^p \\ &\leq \|u_0\|_{H_x^{s_c}} + C(d, p)(2\eta + 2C(d, p)\|u_0\|_{L_x^2}) \|\nabla|^{s_c} u\|_{L_t^{p+2} L_x^{\frac{2d(p+2)}{2(d-2)+dp}}(I \times \mathbb{R}^d)}^p \\ &\leq \|u_0\|_{H_x^{s_c}} + C(d, p)(2\eta + 2C(d, p)\|u_0\|_{L_x^2})(2\eta)^p \end{aligned}$$

and similarly,

$$\begin{aligned} \|\Phi(u)\|_{L_t^{p+2} L_x^{\frac{2d(p+2)}{2(d-2)+dp}}(I \times \mathbb{R}^d)} &\leq C(d, p)\|u_0\|_{L_x^2} + C(d, p)\|F(u)\|_{L_t^{\frac{p+2}{p+1}} L_x^{\frac{2d(p+2)}{2(d+2)+dp}}(I \times \mathbb{R}^d)} \\ &\leq C(d, p)\|u_0\|_{L_x^2} + C(d, p)\|u_0\|_{L_x^2}(2\eta)^p. \end{aligned}$$

Arguing as above and invoking (3.1), we obtain

$$\begin{aligned} \|\nabla|^{s_c} \Phi(u)\|_{L_t^{p+2} L_x^{\frac{2d(p+2)}{2(d-2)+dp}}(I \times \mathbb{R}^d)} &\leq \eta + C(d, p) \|\nabla|^{s_c} F(u)\|_{L_t^{\frac{p+2}{p+1}} L_x^{\frac{2d(p+2)}{2(d+2)+dp}}(I \times \mathbb{R}^d)} \\ &\leq \eta + C(d, p)(2\eta)^{1+p}. \end{aligned}$$

Thus, choosing $\eta_0 = \eta_0(d)$ sufficiently small, we see that for $0 < \eta \leq \eta_0$, the functional Φ maps the set $B_1 \cap B_2$ back to itself. To see that Φ is a contraction, we repeat the computations above to obtain

$$\begin{aligned} \|\Phi(u) - \Phi(v)\|_{L_t^{p+2} L_x^{\frac{2d(p+2)}{2(d-2)+dp}}(I \times \mathbb{R}^d)} &\leq C(d, p)\|F(u) - F(v)\|_{L_t^{\frac{p+2}{p+1}} L_x^{\frac{2d(p+2)}{2(d+2)+dp}}(I \times \mathbb{R}^d)} \\ &\leq C(d, p)(2\eta)^p \|u - v\|_{L_t^{p+2} L_x^{\frac{2d(p+2)}{2(d-2)+dp}}(I \times \mathbb{R}^d)}. \end{aligned}$$

Thus, choosing $\eta_0 = \eta_0(d)$ even smaller (if necessary), we can guarantee that Φ is a contraction on the set $B_1 \cap B_2$. By the contraction mapping theorem, it follows that Φ has a fixed point in $B_1 \cap B_2$. Moreover, noting that Φ maps into $C_t^0 H_x^{s_c}$ (not just $L_t^\infty H_x^{s_c}$), we derive (after one more application of the Strichartz inequality) that the fixed point of Φ is indeed a solution to (1.1).

We now turn our attention to the uniqueness statement. Since uniqueness is a local property, it suffices to study a neighbourhood of $t = 0$. By Definition 1.1 (and the Strichartz inequality), any solution to (1.1) belongs to $B_1 \cap B_2$ on some such neighbourhood. Uniqueness thus follows from uniqueness in the contraction mapping theorem.

The claims (3.3) and (3.4) follow from another application of the Strichartz inequality, as above. $\square$

Next, we will establish a stability theory for (1.1) in the settings of Theorems 1.4 and 1.5. We start with the cubic NLS.

Theorem 3.3 (Stability – the cubic). *Let $d \geq 2$ and let I a compact time interval containing zero and $\tilde{u}$ be an approximate solution to (1.1) on $I \times \mathbb{R}^d$ in the sense that*

$$i\tilde{u}_t = -\Delta\tilde{u} + |\tilde{u}|^2\tilde{u} + e$$

for some function e . Assume that

$$\|\tilde{u}\|_{L_t^\infty \dot{H}_x^{\frac{d-2}{2}}(I \times \mathbb{R}^d)} \leq E \quad (3.5)$$

$$S_I(\tilde{u}) \leq L \quad (3.6)$$

for some positive constants E and L . Let $u_0 \in \dot{H}_x^{\frac{d-2}{2}}$ and assume the smallness conditions

$$\|u_0 - \tilde{u}_0\|_{\dot{H}_x^{\frac{d-2}{2}}} \leq \varepsilon \quad (3.7)$$

$$\| |\nabla|^{\frac{d-2}{2}} e \|_{N^0(I)} \leq \varepsilon \quad (3.8)$$

for some $0 < \varepsilon < \varepsilon_1 = \varepsilon_1(E, L)$. Then, there exists a unique strong solution $u : I \times \mathbb{R}^d \mapsto \mathbb{C}$ to (1.1) with initial data u_0 at time $t = 0$ satisfying

$$S_I(u - \tilde{u}) \leq C(E, L)\varepsilon^{d+2} \quad (3.9)$$

$$\| |\nabla|^{\frac{d-2}{2}} (u - \tilde{u}) \|_{S^0(I)} \leq C(E, L)\varepsilon \quad (3.10)$$

$$\| |\nabla|^{\frac{d-2}{2}} u \|_{S^0(I)} \leq C(E, L). \quad (3.11)$$

Proof. We will prove the theorem under the additional assumption that $u_0 \in L_x^2$, so that we can rely on Theorem 3.1 to guarantee that u exists. This additional assumption can be removed *a posteriori* by the usual limiting argument: approximate u_0 in $\dot{H}_x^{s_c}$ by $\{u_n(0)\}_n \subseteq H_x^{s_c}$ and apply the theorem with $\tilde{u} = u_m$, $u = u_n$, and $e = 0$ to deduce that the sequence of solutions $\{u_n\}_n$ with initial data $\{u_n(0)\}_n$ is Cauchy in critical norms and thus convergent to a solution u with initial data u_0 which obeys $|\nabla|^{s_c} u \in S^0(I)$. Thus, it suffices to prove (3.9) through (3.11) as *a priori* estimates, that is we assume that the solution u exists and obeys $|\nabla|^{s_c} u \in S^0(I)$.

We first prove (3.9) through (3.11) under the stronger hypothesis that

$$\| |\nabla|^{\frac{d-2}{2}} \tilde{u} \|_{L_t^{d+2} L_x^{\frac{2d(d+2)}{d^2+2d-4}}(I \times \mathbb{R}^d)} \leq \delta \quad (3.12)$$

for some small $0 < \delta = \delta(d, p)$.

Let $w := u - \tilde{u}$. Then w satisfies the following initial value problem

$$\begin{cases} iw_t = -\Delta w + F(\tilde{u} + w) - F(\tilde{u}) - e \\ w(0) = u_0 - \tilde{u}_0. \end{cases}$$

For $t \in I$ we define

$$A(t) := \| |\nabla|^{\frac{d-2}{2}} [F(\tilde{u} + w) - F(\tilde{u})] \|_{N^0([0, t])}.$$

By the (fractional) chain rule and (3.12),

$$A(t) \lesssim \| |\nabla|^{\frac{d-2}{2}} w \|_{S^0(I)}^3 + \delta \| |\nabla|^{\frac{d-2}{2}} w \|_{S^0(I)}^2 + \delta^2 \| |\nabla|^{\frac{d-2}{2}} w \|_{S^0(I)}. \quad (3.13)$$

On the other hand, by Strichartz, (3.7), and (3.8), we get

$$\| |\nabla|^{\frac{d-2}{2}} w \|_{S^0(I)} \lesssim \| u_0 - \tilde{u}_0 \|_{\dot{H}_x^{\frac{d-2}{2}}} + A(t) + \| |\nabla|^{\frac{d-2}{2}} e \|_{N^0(I)} \lesssim A(t) + \varepsilon. \quad (3.14)$$

Combining (3.13) and (3.14), we obtain

$$A(t) \lesssim (A(t) + \varepsilon)^3 + \delta(A(t) + \varepsilon)^2 + \delta^2(A(t) + \varepsilon) + \varepsilon.$$

A standard continuity argument then shows that if δ is taken sufficiently small,

$$A(t) \lesssim \varepsilon \quad \text{for any } t \in I,$$

which immediately implies (3.9) through (3.11) via an application of the Strichartz inequality and the triangle inequality.

We now prove (3.9) through (3.11) under the hypothesis (3.6), as opposed to (3.12). We first show that

$$\| |\nabla|^{\frac{d-2}{2}} \tilde{u} \|_{S^0(I)} \leq C(E, L). \quad (3.15)$$

Indeed, by (3.6) we may divide I into $J_0 = J_0(L, \eta)$ subintervals $I_j = [t_j, t_{j+1}]$ such that on each spacetime slab $I_j \times \mathbb{R}^d$

$$\| \tilde{u} \|_{L_{t,x}^{d+2}(I_j \times \mathbb{R}^d)} \leq \eta$$

for a small constant $\eta > 0$ to be chosen in a moment. By the Strichartz inequality combined with Corollary 2.7, (3.5), and (3.8),

$$\begin{aligned} \| |\nabla|^{\frac{d-2}{2}} \tilde{u} \|_{S^0(I_j)} &\lesssim \| \tilde{u}(t_j) \|_{\dot{H}_x^{\frac{d-2}{2}}} + \| |\nabla|^{\frac{d-2}{2}} e \|_{N^0(I_j)} + \| |\nabla|^{\frac{d-2}{2}} F(\tilde{u}) \|_{N^0(I_j)} \\ &\lesssim E + \varepsilon + \| \tilde{u} \|_{L_{t,x}^{d+2}(I_j \times \mathbb{R}^d)}^2 \| |\nabla|^{\frac{d-2}{2}} \tilde{u} \|_{S^0(I_j)} \\ &\lesssim E + \varepsilon + \eta^2 \| |\nabla|^{\frac{d-2}{2}} \tilde{u} \|_{S^0(I_j)}. \end{aligned}$$

Thus, choosing $\eta > 0$ small depending on d and ε_1 sufficiently small depending on E , we obtain

$$\| |\nabla|^{\frac{d-2}{2}} \tilde{u} \|_{S^0(I_j)} \lesssim E.$$

Summing this over all subintervals I_j , we derive (3.15). Thus, we may divide I into $J_1 = J_1(E, L)$ subintervals $I_j = [t_j, t_{j+1}]$ such that on each spacetime slab $I_j \times \mathbb{R}^d$

$$\| |\nabla|^{\frac{d-2}{2}} \tilde{u} \|_{L_t^{d+2} L_x^{\frac{2d(d+2)}{d^2+2d-4}}(I_j \times \mathbb{R}^d)} \leq \delta$$

for some small $\delta = \delta(d, p) > 0$ as appearing in (3.12).

Choosing ε_1 sufficiently small (depending on J_1), we can iterate the argument above to obtain for each $0 \leq j < J_1$ and all $0 < \varepsilon < \varepsilon_1$,

$$\begin{aligned} S_{I_j}(u - \tilde{u}) &\leq C(j) \varepsilon^{d+2} \\ \| |\nabla|^{\frac{d-2}{2}}(u - \tilde{u}) \|_{S^0(I_j)} &\leq C(j) \varepsilon \\ \| |\nabla|^{\frac{d-2}{2}} u \|_{S^0(I_j)} &\leq C(j) E \\ \| |\nabla|^{\frac{d-2}{2}} [F(u) - F(\tilde{u})] \|_{N^0(I_j)} &\leq C(j) \varepsilon, \end{aligned} \quad (3.16)$$

provided we can show

$$\| u(t_j) - \tilde{u}(t_j) \|_{\dot{H}_x^{\frac{d-2}{2}}} \leq C(j-1) \varepsilon \quad (3.17)$$

for each $1 \leq j < J_1$. By the Strichartz inequality and the inductive hypothesis,

$$\begin{aligned} \|u(t_j) - \tilde{u}(t_j)\|_{\dot{H}_x^{\frac{d-2}{2}}} &\lesssim \|u_0 - \tilde{u}_0\|_{\dot{H}_x^{\frac{d-2}{2}}} + \left\| |\nabla|^{\frac{d-2}{2}} e \right\|_{N^0([0, t_j])} \\ &\quad + \left\| |\nabla|^{\frac{d-2}{2}} [F(u) - F(\tilde{u})] \right\|_{N^0([0, t_j])} \\ &\lesssim \varepsilon + \sum_{k=0}^{j-1} C(k)\varepsilon, \end{aligned}$$

which proves (3.17).

Summing the bounds in (3.16) over all subintervals I_j , we derive (3.9) through (3.11). This completes the proof of the theorem. $\square$

We now address the stability question in the setting of Theorem 1.5. We will prove the following result.

Theorem 3.4 (Stability). *Let $d \geq 5$ and assume the critical regularity s_c satisfies (1.5). Let I a compact time interval containing zero and let $\tilde{u}$ be an approximate solution to (1.1) on $I \times \mathbb{R}^d$ in the sense that*

$$i\tilde{u}_t = -\Delta\tilde{u} + F(\tilde{u}) + e$$

for some function e . Assume that

$$\|\tilde{u}\|_{L_t^\infty \dot{H}_x^{s_c}(I \times \mathbb{R}^d)} \leq E \tag{3.18}$$

$$S_I(\tilde{u}) \leq L \tag{3.19}$$

for some positive constants E and L . Let $u_0 \in \dot{H}_x^{s_c}$ and assume the smallness conditions

$$\|u_0 - \tilde{u}_0\|_{\dot{H}_x^{s_c}} \leq \varepsilon \tag{3.20}$$

$$\left\| |\nabla|^{s_c} e \right\|_{N^0(I)} \leq \varepsilon \tag{3.21}$$

for some $0 < \varepsilon < \varepsilon_1 = \varepsilon_1(E, L)$. Then, there exists a unique strong solution $u : I \times \mathbb{R}^d \mapsto \mathbb{C}$ to (1.1) with initial data u_0 at time $t = 0$ satisfying

$$S_I(u - \tilde{u}) \leq C(E, L)\varepsilon^{c_1} \tag{3.22}$$

$$\left\| |\nabla|^{s_c}(u - \tilde{u}) \right\|_{S^0(I)} \leq C(E, L)\varepsilon^{c_2} \tag{3.23}$$

$$\left\| |\nabla|^{s_c} u \right\|_{S^0(I)} \leq C(E, L), \tag{3.24}$$

where c_1, c_2 are positive constants that depend on d, p, E , and L .

Remarks. 1. Theorems 3.3 and 3.4 imply the existence and uniqueness of maximal-lifespan solutions in Theorem 3.1. They also prove that the solutions depend uniformly continuously on the initial data (on bounded sets) in norms which are critical with respect to scaling. As a consequence, one can remove from Theorem 3.1 the assumption that the initial data belongs to L_x^2 , since every $\dot{H}_x^{s_c}$ function is well approximated by $H_x^{s_c}$ functions.

2. Using Theorem 3.1 (without the additional assumption that $u_0 \in L_x^2$, due to the first point above), as well as its proof, one easily derives Theorem 1.6. We omit the standard details.

We now turn to the proof of Theorem 3.4; the argument we present is inspired by the one used in the energy-critical setting [20, 32]; see also [23] for a similar technique in the context of the Klein–Gordon equation. The idea is to work in spaces which are critical with respect to scaling but have a small fractional number of derivatives.

For the remainder of this subsection, for any time interval I we will use the abbreviations

$$\begin{aligned} \|u\|_{X^0(I)} &:= \|u\|_{L_t^{q_0} L_x^{\frac{2r_0 d}{d-r_0 s_c}}(I \times \mathbb{R}^d)} \\ \|u\|_{X(I)} &:= \| |\nabla|^{p/2} u \|_{L_t^{q_0} L_x^{\frac{2r_0 d}{2d-r_0(2s_c-p)}}(I \times \mathbb{R}^d)} \\ \|F\|_{Y(I)} &:= \| |\nabla|^{p/2} F \|_{L_t^{\frac{q_0}{1+p}} L_x^{\frac{2r_0 d}{2(1+p)(d-r_0 s_c)+r_0 p}}(I \times \mathbb{R}^d)}, \end{aligned} \quad (3.25)$$

where $(q_0, r_0) = (\frac{2p(2+p)}{p^2-p(d-2)+4}, \frac{d(2+p)}{d-p+s_c(2+p)})$ is a Schrödinger admissible pair. Note that because of (1.5) we have $2 < r_0 < \frac{d}{s_c}$ and $\frac{p(d+2)}{2} < q_0 < \infty$.

First, we connect the spaces in which the solution to (1.1) is measured to the spaces in which the nonlinearity is measured. As usual, this is done via a Strichartz inequality; we reproduce the standard proof.

Lemma 3.5 (Strichartz estimate). *Let I be a compact time interval containing t_0 . Then*

$$\left\| \int_{t_0}^t e^{i(t-s)\Delta} F(s) ds \right\|_{X(I)} \lesssim \|F\|_{Y(I)}.$$

Proof. As $\frac{2r_0 d}{2d-r_0(2s_c-p)}$ and $\frac{2r_0 d}{2(1+p)(d-r_0 s_c)+r_0 p}$ are dual exponents, the dispersive estimate (2.2) implies

$$\|e^{i(t-s)\Delta} F(s)\|_{L_x^{\frac{2r_0 d}{2d-r_0(2s_c-p)}}} \lesssim |t-s|^{-\frac{p(d-r_0 s_c)}{2r_0}} \|F(s)\|_{L_x^{\frac{2r_0 d}{2(1+p)(d-r_0 s_c)+r_0 p}}}.$$

Using the Hardy-Littlewood-Sobolev inequality and the fact that (q_0, r_0) is a Schrödinger admissible pair, we obtain

$$\left\| \int_{t_0}^t e^{i(t-s)\Delta} F(s) ds \right\|_{L_t^{q_0} L_x^{\frac{2r_0 d}{2d-r_0(2s_c-p)}}(I \times \mathbb{R}^d)} \lesssim \|F\|_{L_t^{\frac{q_0}{1+p}} L_x^{\frac{2r_0 d}{2(1+p)(d-r_0 s_c)+r_0 p}}(I \times \mathbb{R}^d)}.$$

As the differentiation operator $|\nabla|^{p/2}$ commutes with the free evolution, we recover the claim. $\square$

Next we establish some connections between the spaces defined in (3.25) and the usual Strichartz spaces.

Lemma 3.6 (Interpolations). *For any compact time interval I ,*

$$\|u\|_{X^0(I)} \lesssim \|u\|_{X(I)} \lesssim \| |\nabla|^{s_c} u \|_{S^0(I)} \quad (3.26)$$

$$\|u\|_{X(I)} \lesssim \|u\|_{L_{t,x}^{\frac{p(d+2)}{2}}(I \times \mathbb{R}^d)}^{\theta_1} \| |\nabla|^{s_c} u \|_{S^0(I)}^{1-\theta_1} \quad (3.27)$$

$$\|u\|_{L_{t,x}^{\frac{p(d+2)}{2}}(I \times \mathbb{R}^d)} \lesssim \|u\|_{X(I)}^{\theta_2} \| |\nabla|^{s_c} u \|_{S^0(I)}^{1-\theta_2}, \quad (3.28)$$

where $0 < \theta_1, \theta_2 < 1$ depend on d, p .

Proof. A simple application of Sobolev embedding yields (3.26).

Using interpolation, we obtain

$$\|u\|_{X(I)} \lesssim \|u\|_{X^0(I)}^{1-\frac{p}{2s_c}} \| |\nabla|^{s_c} u \|_{L_t^{q_0} L_x^{r_0}(I \times \mathbb{R}^d)}^{\frac{p}{2s_c}} \lesssim \|u\|_{X^0(I)}^{1-\frac{p}{2s_c}} \| |\nabla|^{s_c} u \|_{S^0(I)}^{\frac{p}{2s_c}}.$$

On the other hand, as $\frac{p(d+2)}{2} < q_0 < \infty$, interpolation followed by Sobolev embedding yields

$$\|u\|_{X^0(I)} \lesssim \|u\|_{L_{t,x}^{\frac{p(d+2)}{2}}(I \times \mathbb{R}^d)}^{\frac{p(d+2)}{2q_0}} \|u\|_{L_t^\infty L_x^{\frac{pd}{2}}(I \times \mathbb{R}^d)}^{1-\frac{p(d+2)}{2q_0}} \lesssim \|u\|_{L_{t,x}^{\frac{p(d+2)}{2}}(I \times \mathbb{R}^d)}^{\frac{p(d+2)}{2q_0}} \| |\nabla|^{s_c} u \|_{S^0(I)}^{1-\frac{p(d+2)}{2q_0}}.$$

Putting everything together, we derive (3.27).

We now turn to (3.28); using interpolation once again, we obtain

$$\|u\|_{L_{t,x}^{\frac{p(d+2)}{2}}(I \times \mathbb{R}^d)} \lesssim \|u\|_{X^0(I)}^{\frac{q_0(pd-4)}{p[q_0d-2(d+2)]}} \|u\|_{L_t^{\frac{2(d+2)}{d}} L_x^{\frac{pd(d+2)}{2(d+2)-pd}}(I \times \mathbb{R}^d)}^{1 - \frac{q_0(pd-4)}{p[q_0d-2(d+2)]}}$$

and the claim follows from (3.26) and Sobolev embedding. $\square$

Finally, we derive estimates that will help us control the nonlinearity. The main tools we use in deriving these estimates are the fractional chain rules, Lemmas 2.5 and 2.6.

Lemma 3.7 (Nonlinear estimates). *Let $d \geq 5$ and assume the critical regularity s_c satisfies (1.5). Let I a compact time interval. Then,*

$$\|F(u)\|_{Y(I)} \lesssim \|u\|_{X(I)}^{p+1}, \quad (3.29)$$

$$\begin{aligned} & \|F_z(u+v)w\|_{Y(I)} + \|F_{\bar{z}}(u+v)\bar{w}\|_{Y(I)} \\ & \lesssim \left(\|u\|_{X(I)}^{\frac{p(s_c-1)}{s_c}} \|\nabla|^{s_c} u\|_{S^0(I)}^{\frac{p}{s_c}} + \|v\|_{X(I)}^{\frac{p(s_c-1)}{s_c}} \|\nabla|^{s_c} v\|_{S^0(I)}^{\frac{p}{s_c}} \right) \|w\|_{X(I)}, \end{aligned} \quad (3.30)$$

and

$$\begin{aligned} & \| |\nabla|^{s_c} [F(u+v) - F(u)] \|_{N^0(I)} \\ & \lesssim \| |\nabla|^{s_c} v \|_{S^0(I)} \left[\|v\|_{X^0(I)}^p + \|u\|_{X^0(I)}^{p-1+\frac{1}{s_c}} \|v\|_{X^0(I)}^{1-\frac{1}{s_c}} + (\|u\|_{X^0(I)}^{p-1+\frac{1}{s_c}} + \|v\|_{X^0(I)}^{p-1+\frac{1}{s_c}}) \| |\nabla|^{s_c} u \|_{S^0(I)}^{1-\frac{1}{s_c}} \right] \\ & \quad + \| |\nabla|^{s_c} u \|_{S^0(I)} \left(\|v\|_{X^0(I)}^p + \|u\|_{X^0(I)}^\beta \|v\|_{X^0(I)}^{p-\beta} \right) \\ & \quad + \| |\nabla|^{s_c} u \|_{S^0(I)}^{\frac{1}{s_c}} \| |\nabla|^{s_c} v \|_{S^0(I)}^{1-\frac{1}{s_c}} \|u\|_{X^0(I)}^{1-\frac{1}{s_c}} \|v\|_{X^0(I)}^{p-1+\frac{1}{s_c}} \end{aligned} \quad (3.31)$$

for some $0 < \beta < p$.

Proof. Throughout the proof, all spacetime norms are on $I \times \mathbb{R}^d$.

Applying Lemma 2.5 followed (3.26), we find

$$\|F(u)\|_{Y(I)} \lesssim \|u\|_{X^0(I)}^p \|u\|_{X(I)} \lesssim \|u\|_{X(I)}^{1+p}.$$

This establishes (3.29).

We now turn to (3.30); we only treat the first term on the left-hand side, as the second term can be handled similarly. By Lemma 2.4 followed by (3.26),

$$\begin{aligned} \|F_z(u+v)w\|_{Y(I)} & \lesssim \|F_z(u+v)\|_{L_t^{\frac{q_0}{p}} L_x^{\frac{r_0 d}{p(d-r_0 s_c)}}} \|w\|_{X(I)} \\ & \quad + \| |\nabla|^{\frac{p}{2}} F_z(u+v) \|_{L_t^{\frac{q_0}{p}} L_x^{\frac{2r_0 d}{p(2d-2r_0 s_c+r_0)}}} \|w\|_{X^0(I)} \\ & \lesssim \left(\|u+v\|_{X^0(I)}^p + \| |\nabla|^{\frac{p}{2}} F_z(u+v) \|_{L_t^{\frac{q_0}{p}} L_x^{\frac{2r_0 d}{p(2d-2r_0 s_c+r_0)}}} \right) \|w\|_{X(I)}. \end{aligned}$$

Thus, the claim will follow from (3.26), once we establish

$$\begin{aligned} & \| |\nabla|^{\frac{p}{2}} F_z(u+v) \|_{L_t^{\frac{q_0}{p}} L_x^{\frac{2r_0 d}{p(2d-2r_0 s_c+r_0)}}} \\ & \lesssim \|u\|_{X(I)}^{\frac{p(s_c-1)}{s_c}} \| |\nabla|^{s_c} u \|_{S^0(I)}^{\frac{p}{s_c}} + \|v\|_{X(I)}^{\frac{p(s_c-1)}{s_c}} \| |\nabla|^{s_c} v \|_{S^0(I)}^{\frac{p}{s_c}}. \end{aligned} \quad (3.32)$$

For $p \geq 1$, this follows from Lemma 2.5 and (3.26):

$$\| |\nabla|^{\frac{p}{2}} F_z(u+v) \|_{L_t^{\frac{q_0}{p}} L_x^{\frac{2r_0 d}{p(2d-2r_0 s_c+r_0)}}} \lesssim \|u+v\|_{X^0(I)}^{p-1} \|u+v\|_{X(I)} \lesssim \|u+v\|_{X(I)}^p.$$

To derive (3.32) for $p < 1$, we apply Lemma 2.6 (with $s := p/2$ and $1/2 < \sigma < 1$) followed by Hölder's inequality in the time variable, Sobolev embedding, and interpolation:

$$\begin{aligned} \left\| |\nabla|^{\frac{p}{2}} F_z(u+v) \right\|_{L_t^{\frac{q_0}{p}} L_x^{\frac{2r_0 d}{p(2d-2r_0 s_c+r_0)}}} &\lesssim \|u+v\|_{X^0(I)}^{p-\frac{p}{2\sigma}} \left\| |\nabla|^\sigma(u+v) \right\|_{L_t^{q_0} L_x^{\frac{r_0 d}{d-r_0(s_c-\sigma)}}}^{\frac{p}{2\sigma}} \\ &\lesssim \left\| |\nabla|^\sigma(u+v) \right\|_{L_t^{q_0} L_x^{\frac{r_0 d}{d-r_0(s_c-\sigma)}}}^p \\ &\lesssim \|u\|_{X^0(I)}^{p-\frac{p\sigma}{s_c}} \left\| |\nabla|^{s_c} u \right\|_{S^0(I)}^{\frac{p\sigma}{s_c}} + \|v\|_{X^0(I)}^{p-\frac{p\sigma}{s_c}} \left\| |\nabla|^{s_c} v \right\|_{S^0(I)}^{\frac{p\sigma}{s_c}}. \end{aligned}$$

Invoking (3.26), this settles (3.32) and hence (3.30).

To prove (3.31), we estimate

$$\begin{aligned} \left\| |\nabla|^{s_c} [F(u+v) - F(u)] \right\|_{N^0(I)} &\lesssim \left\| |\nabla|^{s_c-1} [\nabla v \cdot F'(u+v)] \right\|_{N^0(I)} \\ &\quad + \left\| |\nabla|^{s_c-1} [\nabla u \cdot (F'(u+v) - F'(u))] \right\|_{N^0(I)}. \end{aligned} \quad (3.33)$$

To estimate the first term on the right-hand side of (3.33), we use Lemmas 2.4 and 2.6 together with Hölder's inequality and interpolation:

$$\begin{aligned} &\left\| |\nabla|^{s_c-1} [\nabla v \cdot F'(u+v)] \right\|_{N^0(I)} \\ &\lesssim \left\| |\nabla|^{s_c} v \right\|_{S^0(I)} (\|u\|_{X^0(I)}^p + \|v\|_{X^0(I)}^p) \\ &\quad + \left\| |\nabla|^{s_c} v \right\|_{S^0(I)}^{\frac{1}{s_c}} \|v\|_{X^0(I)}^{1-\frac{1}{s_c}} \|u+v\|_{X^0(I)}^{p-1+\frac{1}{s_c}} \left\| |\nabla|^{s_c}(u+v) \right\|_{S^0(I)}^{1-\frac{1}{s_c}} \\ &\lesssim \left\| |\nabla|^{s_c} v \right\|_{S^0(I)} \left[\|v\|_{X^0(I)}^p + \|u\|_{X^0(I)}^{p-1+\frac{1}{s_c}} \|v\|_{X^0(I)}^{1-\frac{1}{s_c}} \right. \\ &\quad \left. + (\|u\|_{X^0(I)}^{p-1+\frac{1}{s_c}} + \|v\|_{X^0(I)}^{p-1+\frac{1}{s_c}}) \left\| |\nabla|^{s_c} u \right\|_{S^0(I)}^{1-\frac{1}{s_c}} \right]. \end{aligned}$$

To estimate the second term on the right-hand side of (3.33), we use Lemma 2.8 together with Hölder's inequality, interpolation, and (3.26):

$$\begin{aligned} &\left\| |\nabla|^{s_c-1} [\nabla u \cdot (F'(u+v) - F'(u))] \right\|_{N^0(I)} \\ &\lesssim \left\| |\nabla|^{s_c} u \right\|_{S^0(I)} \|v\|_{X^0(I)}^p + \left\| |\nabla|^{s_c} u \right\|_{S^0(I)} \|u\|_{X^0(I)}^{\frac{s_c-1}{\sigma}} \|v\|_{X^0(I)}^{p-\frac{s_c-1}{\sigma}} \\ &\quad + \left\| |\nabla|^{s_c} u \right\|_{S^0(I)}^{\frac{1}{s_c}} \|u\|_{X^0(I)}^{1-\frac{1}{s_c}} \left\| |\nabla|^{s_c} v \right\|_{S^0(I)}^{1-\frac{1}{s_c}} \|v\|_{X^0(I)}^{p-1+\frac{1}{s_c}}, \end{aligned}$$

where $s_c - 1 < \sigma p < p$. Denoting $\beta := \frac{s_c-1}{\sigma}$ and collecting all the estimates above we derive (3.31). $\square$

We have now all the tools we need to attack Theorem 3.4. We start with the following:

Lemma 3.8 (Short-time perturbations). *Let $d \geq 5$ and assume the critical regularity s_c satisfies (1.5). Let I be a compact time interval containing zero and let $\tilde{u}$ be an approximate solution to (1.1) on $I \times \mathbb{R}^d$ in the sense that*

$$i\tilde{u}_t = -\Delta\tilde{u} + F(\tilde{u}) + e$$

for some function e . Assume that

$$\|\tilde{u}\|_{L_t^\infty \dot{H}_x^{s_c}(I \times \mathbb{R}^d)} \leq E$$

for some positive constant E . Moreover, let $u_0 \in \dot{H}_x^{s_c}$ and assume that

$$\|\tilde{u}\|_{X(I)} \leq \delta \quad (3.34)$$

$$\|u_0 - \tilde{u}_0\|_{\dot{H}_x^{s_c}} \leq \varepsilon \quad (3.35)$$

$$\left\| |\nabla|^{s_c} e \right\|_{N^0(I)} \leq \varepsilon \quad (3.36)$$

for some small $0 < \delta = \delta(E)$ and $0 < \varepsilon < \varepsilon_0(E)$. Then there exists a unique solution $u : I \times \mathbb{R}^d \rightarrow \mathbb{C}$ to (1.1) with initial data u_0 at time $t = 0$; it satisfies

$$\|u - \tilde{u}\|_{X(I)} \lesssim \varepsilon \quad (3.37)$$

$$\| |\nabla|^{s_c}(u - \tilde{u}) \|_{S^0(I)} \lesssim \varepsilon^{c(d,p)} \quad (3.38)$$

$$\| |\nabla|^{s_c} u \|_{S^0(I)} \lesssim E \quad (3.39)$$

$$\|F(u) - F(\tilde{u})\|_{Y(I)} \lesssim \varepsilon \quad (3.40)$$

$$\| |\nabla|^{s_c} [F(u) - F(\tilde{u})] \|_{N^0(I)} \lesssim \varepsilon^{c(d,p)}, \quad (3.41)$$

for some positive constant $c(d, p)$.

Proof. As explained at the beginning of the proof of Theorem 3.3, we may assume that u exists and merely show that it obeys the estimates stated above.

We start by deriving some bounds on $\tilde{u}$ and u . By Strichartz, Corollary 2.7, Lemma 3.6, (3.34), and (3.36),

$$\begin{aligned} \| |\nabla|^{s_c} \tilde{u} \|_{S^0(I)} &\lesssim \|\tilde{u}\|_{L_t^\infty \dot{H}_x^{s_c}(I \times \mathbb{R}^d)} + \| |\nabla|^{s_c} F(\tilde{u}) \|_{N^0(I)} + \| |\nabla|^{s_c} e \|_{N^0(I)} \\ &\lesssim E + \|\tilde{u}\|_{L_{t,x}^{\frac{p(d+2)}{2}}(I \times \mathbb{R}^d)}^p \| |\nabla|^{s_c} \tilde{u} \|_{S^0(I)} + \varepsilon \\ &\lesssim E + \delta^{p\theta_2} \| |\nabla|^{s_c} \tilde{u} \|_{S^0(I)}^{1+p(1-\theta_2)} + \varepsilon, \end{aligned}$$

where θ_2 is as in Lemma 3.6. Choosing δ small depending on d, p, E and ε_0 sufficiently small depending on E , we obtain

$$\| |\nabla|^{s_c} \tilde{u} \|_{S^0(I)} \lesssim E. \quad (3.42)$$

Moreover, by Lemma 3.5, Lemma 3.7, (3.34), and (3.36),

$$\|e^{it\Delta} \tilde{u}_0\|_{X(I)} \lesssim \|\tilde{u}\|_{X(I)} + \|F(\tilde{u})\|_{Y(I)} + \| |\nabla|^{s_c} e \|_{N^0(I)} \lesssim \delta + \delta^{\frac{d+2}{d-2}} + \varepsilon \lesssim \delta,$$

provided δ and ε_0 are chosen sufficiently small. Combining this with the triangle inequality, (3.26), the Strichartz inequality, and (3.35), we obtain

$$\|e^{it\Delta} u_0\|_{X(I)} \lesssim \|e^{it\Delta} \tilde{u}_0\|_{X(I)} + \|u_0 - \tilde{u}_0\|_{\dot{H}_x^{s_c}} \lesssim \delta + \varepsilon \lesssim \delta.$$

Thus, another application of Lemma 3.5 combined with Lemma 3.7 gives

$$\|u\|_{X(I)} \lesssim \|e^{it\Delta} u_0\|_{X(I)} + \|F(u)\|_{Y(I)} \lesssim \delta + \|u\|_{X(I)}^{\frac{d+2}{d-2}}.$$

Choosing δ sufficiently small, the usual bootstrap argument yields

$$\|u\|_{X(I)} \lesssim \delta. \quad (3.43)$$

Next we derive the claimed bounds on $w := u - \tilde{u}$. Note that w is a solution to

$$\begin{cases} iw_t = -\Delta w + F(\tilde{u} + w) - F(\tilde{u}) - e \\ w(t_0) = u_0 - \tilde{u}_0. \end{cases}$$

Using Lemma 3.5 together with Lemma 3.6, the Strichartz inequality, (3.35), and (3.36), we see that

$$\|w\|_{X(I)} \lesssim \|u_0 - \tilde{u}_0\|_{\dot{H}_x^{s_c}} + \| |\nabla|^{s_c} e \|_{N^0(I)} + \|F(u) - F(\tilde{u})\|_{Y(I)} \lesssim \varepsilon + \|F(u) - F(\tilde{u})\|_{Y(I)}.$$

To estimate the difference of the nonlinearities, we use Lemma 3.7, (3.34), (3.42):

$$\begin{aligned} \|F(u) - F(\tilde{u})\|_{Y(I)} &\lesssim \left[\|\tilde{u}\|_{X(I)}^{\frac{p(s_c-1)}{s_c}} \| |\nabla|^{s_c} \tilde{u} \|_{S^0(I)}^{\frac{p}{s_c}} + \|w\|_{X(I)}^{\frac{p(s_c-1)}{s_c}} \| |\nabla|^{s_c} w \|_{S^0(I)}^{\frac{p}{s_c}} \right] \|w\|_{X(I)} \\ &\lesssim \delta^{\frac{p(s_c-1)}{s_c}} E^{\frac{p}{s_c}} \|w\|_{X(I)} + \| |\nabla|^{s_c} w \|_{S^0(I)}^{\frac{p}{s_c}} \|w\|_{X(I)}^{1+\frac{p(s_c-1)}{s_c}}. \end{aligned} \quad (3.44)$$

Thus, choosing δ sufficiently small depending only on E , we obtain

$$\|w\|_{X(I)} \lesssim \varepsilon + \| |\nabla|^{s_c} w \|_{S^0(I)}^{\frac{p}{s_c}} \|w\|_{X(I)}^{1+\frac{p(s_c-1)}{s_c}}. \quad (3.45)$$

On the other hand, by the Strichartz inequality and the hypotheses,

$$\begin{aligned} \| |\nabla|^{s_c} w \|_{S^0(I)} &\lesssim \|u_0 - \tilde{u}_0\|_{\dot{H}_x^{s_c}} + \| |\nabla|^{s_c} e \|_{N^0(I)} + \| |\nabla|^{s_c} [F(u) - F(\tilde{u})] \|_{N^0(I)} \\ &\lesssim \varepsilon + \| |\nabla|^{s_c} [F(u) - F(\tilde{u})] \|_{N^0(I)}. \end{aligned} \quad (3.46)$$

To estimate the difference of the nonlinearities, we use (3.31) together with Lemma 3.6, (3.34), (3.42), and (3.43),

$$\begin{aligned} \| |\nabla|^{s_c} [F(u) - F(\tilde{u})] \|_{N^0(I)} &\lesssim \| |\nabla|^{s_c} w \|_{S^0(I)} \left(\delta + \delta^{p-1+\frac{1}{s_c}} E^{1-\frac{1}{s_c}} \right) \\ &\quad + E \delta^\beta \|w\|_{X(I)}^{p-\beta} + \delta^{1-\frac{1}{s_c}} E^{\frac{1}{s_c}} \| |\nabla|^{s_c} w \|_{S^0(I)}^{1-\frac{1}{s_c}} \|w\|_{X(I)}^{p-1+\frac{1}{s_c}} \end{aligned} \quad (3.47)$$

for some $0 < \beta < p$. Thus, choosing δ small depending only on E , (3.46) implies

$$\| |\nabla|^{s_c} w \|_{S^0(I)} \lesssim \varepsilon + \|w\|_{X(I)}^{p-\beta} + \| |\nabla|^{s_c} w \|_{S^0(I)}^{1-\frac{1}{s_c}} \|w\|_{X(I)}^{p-1+\frac{1}{s_c}}. \quad (3.48)$$

Combining (3.45) with (3.48), the usual bootstrap argument yields (3.37) and (3.38), provided ε_0 is chosen sufficiently small depending on E . By the triangle inequality, (3.38) and (3.42) imply (3.39).

Claims (3.40) and (3.41) follow from (3.44) and (3.47) combined with (3.37) and (3.38), provided we take δ and ε_0 sufficiently small depending on E . $\square$

We are finally in a position to prove the stability result.

Proof of Theorem 3.4. Our first goal is to show

$$\| |\nabla|^{s_c} \tilde{u} \|_{S^0(I)} \leq C(E, L). \quad (3.49)$$

Indeed, by (3.19) we may divide I into $J_0 = J_0(L, \eta)$ subintervals $I_j = [t_j, t_{j+1}]$ such that on each spacetime slab $I_j \times \mathbb{R}^d$

$$\|\tilde{u}\|_{L_{t,x}^{\frac{p(d+2)}{2}}(I_j \times \mathbb{R}^d)} \leq \eta$$

for a small constant $\eta > 0$ to be chosen in a moment. By the Strichartz inequality combined with Corollary 2.7, (3.18) and (3.21),

$$\begin{aligned} \| |\nabla|^{s_c} \tilde{u} \|_{S^0(I_j)} &\lesssim \|\tilde{u}(t_j)\|_{\dot{H}_x^{s_c}} + \| |\nabla|^{s_c} e \|_{N^0(I_j)} + \| |\nabla|^{s_c} F(\tilde{u}) \|_{N^0(I_j)} \\ &\lesssim E + \varepsilon + \|\tilde{u}\|_{L_{t,x}^{\frac{p(d+2)}{2}}(I_j \times \mathbb{R}^d)}^p \| |\nabla|^{s_c} \tilde{u} \|_{S^0(I_j)} \\ &\lesssim E + \varepsilon + \eta^p \| |\nabla|^{s_c} \tilde{u} \|_{S^0(I_j)}. \end{aligned}$$

Thus, choosing $\eta > 0$ small depending on d, p and ε_1 sufficiently small depending on E , we obtain

$$\| |\nabla|^{s_c} \tilde{u} \|_{S^0(I_j)} \lesssim E.$$

Summing this over all subintervals I_j , we derive (3.49).

Using Lemma 3.6 together with (3.49), we obtain

$$\|\tilde{u}\|_{X(I)} \leq C(E, L) \quad (3.50)$$

By (3.50), we may divide I into $J_1 = J_1(E, L)$ subintervals $I_j = [t_j, t_{j+1}]$ such that on each spacetime slab $I_j \times \mathbb{R}^d$

$$\|\tilde{u}\|_{X(I_j)} \leq \delta$$

for some small $\delta = \delta(E) > 0$ as in Lemma 3.8.

Thus, choosing ε_1 sufficiently small (depending on J_1 and E), we may apply Lemma 3.8 to obtain for each $0 \leq j < J_1$ and all $0 < \varepsilon < \varepsilon_1$,

$$\begin{aligned} \|u - \tilde{u}\|_{X(I_j)} &\leq C(j)\varepsilon^{c(d,p)^j} \\ \||\nabla|^{s_c}(u - \tilde{u})\|_{S^0(I_j)} &\leq C(j)\varepsilon^{c(d,p)^{j+1}} \\ \||\nabla|^{s_c}u\|_{\dot{S}^1(I_j)} &\leq C(j)E \\ \|F(u) - F(\tilde{u})\|_{Y(I_j)} &\leq C(j)\varepsilon^{c(d,p)^j} \\ \||\nabla|^{s_c}[F(u) - F(\tilde{u})]\|_{N^0(I_j)} &\leq C(j)\varepsilon^{c(d,p)^{j+1}}, \end{aligned} \tag{3.51}$$

provided we can show

$$\|u(t_j) - \tilde{u}(t_j)\|_{\dot{H}^{s_c}} \lesssim \varepsilon^{c(d,p)^j} \leq \varepsilon_0 \tag{3.52}$$

for each $0 \leq j < J_1$, where ε_0 is as in Lemma 3.8. By the Strichartz inequality and the inductive hypothesis,

$$\begin{aligned} \|u(t_j) - \tilde{u}(t_j)\|_{\dot{H}^{s_c}} &\lesssim \|u_0 - \tilde{u}_0\|_{\dot{H}^{s_c}} + \||\nabla|^{s_c}e\|_{N^0([0,t_j])} + \||\nabla|^{s_c}[F(u) - F(\tilde{u})]\|_{N^0([0,t_j])} \\ &\lesssim \varepsilon + \sum_{k=0}^{j-1} C(k)\varepsilon^{c(d,p)^{k+1}}. \end{aligned}$$

Taking ε_1 sufficiently small compared to ε_0 , we see that (3.52) is satisfied.

Summing the bounds in (3.51) over all subintervals I_j and using Lemma 3.6, we derive (3.22) through (3.24). This completes the proof of the theorem. $\square$

4. REDUCTION TO ALMOST PERIODIC SOLUTIONS

The goal of this section is to prove Theorem 1.12. In order to achieve this, we repeat the argument presented in [19]. Since the procedure is by now standard, we content ourselves with identifying the main steps and indicating, whenever necessary, the changes that appear with respect to the presentation in [19]. See also [13, 14, 15, 20, 34] for similar arguments in other contexts.

We start by presenting the setup. Throughout this section, we fix a dimension $d \geq 5$ and assume that either the critical regularity is $s_c = \frac{d-2}{2}$ or it satisfies (1.5). For any $0 \leq E_0 < \infty$, we define

$$L(E_0) := \sup\{S_I(u) : u : I \times \mathbb{R}^d \rightarrow \mathbb{C} \text{ such that } \sup_{t \in I} \||\nabla|^{s_c}u(t)\|_2^2 \leq E_0\},$$

where the supremum is taken over all solutions $u : I \times \mathbb{R}^d \rightarrow \mathbb{C}$ to (1.1) obeying $\||\nabla|^{s_c}u(t)\|_2^2 \leq E_0$. Thus, $L : [0, \infty) \rightarrow [0, \infty]$ is a non-decreasing function. Moreover, from Theorem 1.6,

$$L(E_0) \lesssim_d E_0^{\frac{p(d+2)}{4}} \quad \text{for } E_0 \leq \eta_0,$$

where $\eta_0 = \eta_0(d, p)$ is the threshold from the small data theory.

From Theorems 3.3 and 3.4, we see that L is continuous in both settings considered here. Therefore, there must exist a unique *critical* $E_c \in (0, \infty]$ such that $L(E_0) < \infty$ for $E_0 < E_c$ and $L(E_0) = \infty$ for $E_0 \geq E_c$. In particular, if $u : I \times \mathbb{R}^d \rightarrow \mathbb{C}$ is a maximal-lifespan solution to (1.1) such that $\sup_{t \in I} \||\nabla|^{s_c}u(t)\|_2^2 < E_c$, then u is global and moreover,

$$S_{\mathbb{R}}(u) \leq L(\|u\|_{L_t^\infty \dot{H}_x^{s_c}}^2).$$

Failure of Theorems 1.4 or 1.5 is equivalent to $0 < E_c < \infty$.

Following the presentation in [19, 34], the main step in proving Theorem 1.12 is to prove a Palais–Smale condition modulo the symmetries of the equation. With the Palais–Smale condition in place, the proof of Theorem 1.12 is standard; see, for example, [13, 14, 19, 34].

Proposition 4.1 (Palais-Smale condition modulo symmetries). *Assume we are either in the setting of Theorem 1.4 or Theorem 1.5. Let $u_n : I_n \times \mathbb{R}^d \mapsto \mathbb{C}$ be a sequence of solutions to (1.1) such that*

$$\limsup_{n \rightarrow \infty} \sup_{t \in I_n} \|\nabla |^{s_c} u_n(t)\|_2^2 = E_c \quad (4.1)$$

and let $t_n \in I_n$ be a sequence of times such that

$$\lim_{n \rightarrow \infty} S_{\geq t_n}(u_n) = \lim_{n \rightarrow \infty} S_{\leq t_n}(u_n) = \infty.$$

Then the sequence $u_n(t_n)$ has a subsequence which converges in $\dot{H}_x^{s_c}(\mathbb{R}^d)$ modulo symmetries.

The proof of this proposition follows the same recipe as that used to prove the analogous statement in [19]. The main ingredients are the linear profile decomposition from Lemma 2.12 and the stability results Theorems 3.3 and 3.4. The only new difficulty appears when one endeavors to prove decoupling of the nonlinear profiles. In the energy-critical setting, one uses the pointwise inequality

$$\left| \nabla \left(\sum_{j=1}^J F(f_j) - F\left(\sum_{j=1}^J f_j\right) \right) \right| \lesssim_J \sum_{j \neq j'} |\nabla f_j| |f_{j'}|^{\frac{4}{d-2}},$$

which does not carry over to the case of a non-integer number of derivatives. Let us first discuss the case $0 < p \leq 1$; in this scenario, we use (2.6) to obtain the following substitute for the inequality above:

$$\begin{aligned} & \mathcal{D}_{s_c-1} \left(\nabla \left[\sum_{j=1}^J F(f_j) - F\left(\sum_{j=1}^J f_j\right) \right] \right) \\ & \lesssim \mathcal{D}_{s_c-1} \left(\sum_{j=1}^J \nabla f_j \left[F'(f_j) - F'\left(\sum_{j'=1}^J f_{j'}\right) \right] \right) \\ & \lesssim_J \sum_{j \neq j'} \left\{ \mathcal{D}_{s_c-1}(\nabla f_j) |f_{j'}|^p + \left[\sum_{l=1}^J \mathcal{D}_\sigma(f_l) \right]^{\frac{\sigma}{\sigma-1}} M(|\nabla f_j|^{\frac{1}{1-p}})^{1-p} M(f_{j'})^{p-\frac{\sigma}{\sigma-1}} \right\} \end{aligned}$$

for some $s_c - 1 < \sigma p < p$. We remind the reader that M denotes the Hardy–Littlewood maximal function, which commutes with the symmetries of the equation. The operator $\mathcal{D}_s$ is defined in (2.5) and it behaves like $|\nabla|^s$ under symmetries.

In the case discussed above, one has both a small-power non-polynomial nonlinearity and a non-integer number of derivatives $1 < s_c < 2$, which makes it the most awkward of the scenarios we need to consider. The remaining cases of Theorems 1.4 and 1.5, can be handled using various permutations of the techniques discussed above or some alternatives. In particular, we draw the reader's attention to [14] which considers the cubic nonlinearity with $s_c = \frac{1}{2}$.

5. THE FINITE-TIME BLOWUP SOLUTION

In this section we preclude scenario I described in Theorem 1.13. We start by considering finite-time blowup solutions in the setting of Theorem 1.5.

Theorem 5.1 (Absence of finite-time blowup solutions). *Let $d \geq 5$ and assume the critical regularity s_c satisfies (1.5). Then there are no finite-time blowup solutions to (1.1) in the sense of Theorem 1.13.*

Proof. We argue by contradiction. Assume that there exists a solution $u : I \times \mathbb{R}^d \rightarrow \mathbb{C}$ that is a finite-time blowup solution in the sense of Theorem 1.13. Assume also, without loss of generality, that the solution u blows up in finite time in the future, that is, $T := \sup I < \infty$.

By hypothesis and Sobolev embedding,

$$\|u\|_{L_t^\infty L_x^{\frac{dp}{2}}(I \times \mathbb{R}^d)} \lesssim \| |\nabla|^{s_c-1} u \|_{L_t^\infty L_x^{\frac{2d}{d-2}}(I \times \mathbb{R}^d)} \lesssim \|u\|_{L_t^\infty \dot{H}_x^{s_c}(I \times \mathbb{R}^d)} \lesssim_u 1.$$

Thus, using the Duhamel formula (1.7) into the future together with the Strichartz and Hölder inequalities, as well as the fractional chain rule, we obtain

$$\begin{aligned} \| |\nabla|^{s_c-1} u(t) \|_2 &\leq \left\| \int_t^T e^{i(t-s)\Delta} |\nabla|^{s_c-1} F(u(s)) ds \right\|_2 \\ &\lesssim \| |\nabla|^{s_c-1} F(u) \|_{L_t^2 L_x^{\frac{2d}{d+2}}([t, T] \times \mathbb{R}^d)} \\ &\lesssim (T-t)^{\frac{1}{2}} \| |\nabla|^{s_c-1} u \|_{L_t^\infty L_x^{\frac{2d}{d-2}}([t, T] \times \mathbb{R}^d)} \|u\|_{L_t^\infty L_x^{\frac{dp}{2}}([t, T] \times \mathbb{R}^d)}^p \\ &\lesssim_u (T-t)^{\frac{1}{2}}. \end{aligned}$$

Interpolating with $u \in L_t^\infty \dot{H}_x^{s_c}$ (recalling that $1 < s_c < 2$ by (1.5)) we derive that the energy $E(u) \rightarrow 0$ as $t \rightarrow T$. Invoking the conservation of energy, we deduce that $u \equiv 0$. This contradicts the fact that u is a blowup solution. $\square$

We consider next finite-time blowup solutions in the setting of Theorem 1.4.

Theorem 5.2 (Absence of finite-time blowup solutions – the cubic). *Let $d \geq 5$ and assume $p = 2$. Then there are no finite-time blowup solutions to (1.1) in the sense of Theorem 1.13.*

Proof. Again, we argue by contradiction. Let $u : I \times \mathbb{R}^d \rightarrow \mathbb{C}$ be a finite-time blowup solution in the sense of Theorem 1.13 and assume that $T := \sup I < \infty$.

By Sobolev embedding and the hypothesis,

$$\|u\|_{L_t^\infty L_x^d(I \times \mathbb{R}^d)} \lesssim \| |\nabla|^{\frac{d-4}{2}} u \|_{L_t^\infty L_x^{\frac{2d}{d-2}}(I \times \mathbb{R}^d)} \lesssim \|u\|_{L_t^\infty \dot{H}_x^{\frac{d-2}{2}}(I \times \mathbb{R}^d)} \lesssim_u 1.$$

Thus, using the Duhamel formula (1.7) into the future together with the Strichartz and Hölder inequalities, as well as the fractional chain rule, we obtain

$$\begin{aligned} \| |\nabla|^{\frac{d-4}{2}} u(t) \|_2 &\leq \left\| \int_t^T e^{i(t-s)\Delta} |\nabla|^{\frac{d-4}{2}} F(u(s)) ds \right\|_2 \\ &\lesssim \| |\nabla|^{\frac{d-4}{2}} F(u) \|_{L_t^2 L_x^{\frac{2d}{d+2}}([t, T] \times \mathbb{R}^d)} \\ &\lesssim (T-t)^{\frac{1}{2}} \| |\nabla|^{\frac{d-4}{2}} u \|_{L_t^\infty L_x^{\frac{2d}{d-2}}([t, T] \times \mathbb{R}^d)} \|u\|_{L_t^\infty L_x^d([t, T] \times \mathbb{R}^d)}^2 \\ &\lesssim_u (T-t)^{\frac{1}{2}}. \end{aligned} \tag{5.1}$$

In particular, $u(t) \in \dot{H}_x^{\frac{d-4}{2}}$.

In the case when $d = 5$, one can interpolate between (5.1) and the hypothesis $u \in L_t^\infty \dot{H}_x^{\frac{d-2}{2}}(I \times \mathbb{R}^d)$ to derive that $\| \nabla u(t) \|_2 \rightarrow 0$ as $t \rightarrow T$, and hence, by Sobolev embedding, the energy $E(u(t)) \rightarrow 0$ as $t \rightarrow T$. Using the conservation of energy, we deduce that $u \equiv 0$. This contradicts the fact that for a finite-time blowup solution, $S_I(u) = \infty$.

To handle higher dimensions, we iterate the computations in (5.1) with one less derivative to deduce that

$$\| |\nabla|^{\frac{d-6}{2}} u(t) \|_2 \lesssim_u (T-t).$$

For $d = 6$ this immediately implies that u must have zero mass, while for $d = 7$, the argument used to handle dimension $d = 5$ implies that u must have zero energy. In both cases, we derive a contradiction to the fact that u is a blowup solution.

To derive a contradiction for dimensions $d \geq 8$, we iterate the argument presented above.

This finishes the proof of the theorem. $\square$

6. NEGATIVE REGULARITY

In this section we prove that in scenarios II and III described in Theorem 1.13, the solution u admits negative regularity; more precisely, it lies in $L_t^\infty \dot{H}_x^{-\varepsilon}$ for some $\varepsilon > 0$. In particular, this shows that the solution decays sufficiently rapidly (in space) to belong to $L_t^\infty L_x^2$. We first consider the setting of Theorem 1.5. At the end of this section we explain the changes needed to prove negative regularity in the setting of Theorem 1.4.

Theorem 6.1 (Negative regularity for scenarios II and III). *Let $d \geq 5$ and assume the critical regularity s_c obeys (1.5). Let u be a global solution to (1.1) that is almost periodic modulo symmetries. Suppose also that*

$$\sup_{t \in I} \|\nabla|^{s_c} u(t)\|_{L_x^2} < \infty \quad (6.1)$$

and

$$\inf_{t \in I} N(t) \geq 1. \quad (6.2)$$

Then $u \in L_t^\infty \dot{H}_x^{-\varepsilon}$ for some $\varepsilon = \varepsilon(d) > 0$. In particular, $u \in L_t^\infty L_x^2$.

To prove Theorem 6.1, we employ the argument used in [19] to treat the energy-critical case, $s_c = 1$. We achieve our goal in two steps: First, we ‘break’ scaling in a Lebesgue space; more precisely, we prove that our solution lives in $L_t^\infty L_x^q$ for some $2 < q < \frac{dp}{2}$. Next, we use a double Duhamel trick to upgrade this to $u \in L_t^\infty \dot{H}_x^{s_c - s_0}$ for some $s_0 = s_0(d, p, q) > 0$. Iterating the second step finitely many times, we arrive at Theorem 6.1.

Let u be a solution to (1.1) that obeys the hypotheses of Theorem 6.1. Let $\eta > 0$ be a small constant to be chosen later. Then by Remark 1.11 combined with (6.2), there exists $N_0 = N_0(\eta)$ such that

$$\|\nabla|^{s_c} u_{\leq N_0}\|_{L_t^\infty L_x^2} \leq \eta. \quad (6.3)$$

We turn now to our first step, that is, breaking scaling in a Lebesgue space. To this end, we define

$$A(N) := \begin{cases} N^{\frac{d-3}{2} - \frac{2}{p}} \sup_{t \in \mathbb{R}} \|u_N(t)\|_{L_x^{\frac{2d}{d-3}}} & \text{for } d = 5, 6 \\ N^{\frac{d-2}{2} - \frac{2}{p} - \frac{p}{2}} \sup_{t \in \mathbb{R}} \|u_N(t)\|_{L_x^{\frac{2d}{d-2-p}}} & \text{for } d \geq 7 \end{cases}$$

for frequencies $N \leq 10N_0$. To simplify the formulas appearing below, we introduce the notation

$$\alpha(d) := \begin{cases} -\frac{d-3}{2} + \frac{2}{p} & \text{for } d = 5, 6 \\ -\frac{d-2}{2} + \frac{2}{p} + \frac{p}{2} & \text{for } d \geq 7. \end{cases}$$

Note that by (1.5), $0 < \alpha(d) < 1$ for $d = 5, 6$ and $0 < \alpha(d) < p$ for $d \geq 7$. Note also that by Bernstein’s inequality and (6.1),

$$A(N) \lesssim N^{\frac{d}{2} - \frac{2}{p}} \|u_N\|_{L_t^\infty L_x^2} \lesssim \|\nabla|^{s_c} u\|_{L_t^\infty L_x^2} < \infty. \quad (6.4)$$

We next prove a recurrence formula for $A(N)$.

Lemma 6.2 (Recurrence). *For all $N \leq 10N_0$,*

$$A(N) \lesssim_u \left(\frac{N}{N_0}\right)^{\min\{1, p\} - \alpha(d)} + \eta^p \sum_{\frac{N}{10} \leq N_1 \leq N_0} \left(\frac{N}{N_1}\right)^{\min\{1, p\} - \alpha(d)} A(N_1) + \eta^p \sum_{N_1 < \frac{N}{10}} \left(\frac{N_1}{N}\right)^{\alpha(d)} A(N_1).$$

Proof. We first give the proof in dimensions $d \geq 7$. Once this is completed, we will explain the changes necessary to treat $d = 5, 6$.

Fix $N \leq 10N_0$. By time-translation symmetry, it suffices to prove

$$\begin{aligned} N^{-\alpha(d)} \|u_N(0)\|_{L_x^{\frac{2d}{d-2-p}}} &\lesssim_u \left(\frac{N}{N_0}\right)^{p-\alpha(d)} + \eta^p \sum_{\frac{N}{10} \leq N_1 \leq N_0} \left(\frac{N}{N_1}\right)^{p-\alpha(d)} A(N_1) \\ &\quad + \eta^p \sum_{N_1 < \frac{N}{10}} \left(\frac{N_1}{N}\right)^{\alpha(d)} A(N_1). \end{aligned} \quad (6.5)$$

Using the Duhamel formula (1.7) into the future followed by the triangle inequality, Bernstein, and the dispersive inequality, we estimate

$$\begin{aligned} N^{-\alpha(d)} \|u_N(0)\|_{L_x^{\frac{2d}{d-2-p}}} &\leq N^{-\alpha(d)} \int_0^{N^{-2}} \|e^{-it\Delta} P_N F(u(t))\|_{L_x^{\frac{2d}{d-2-p}}} dt \\ &\quad + N^{-\alpha(d)} \int_{N^{-2}}^\infty \|e^{-it\Delta} P_N F(u(t))\|_{L_x^{\frac{2d}{d-2-p}}} dt \\ &\lesssim N^{\frac{p}{2}+1-\alpha(d)} \int_0^{N^{-2}} \|e^{-it\Delta} P_N F(u(t))\|_{L_x^2} dt \\ &\quad + N^{-\alpha(d)} \|P_N F(u)\|_{L_t^\infty L_x^{\frac{2d}{d+2+p}}} \int_{N^{-2}}^\infty t^{-\frac{2+p}{2}} dt \\ &\lesssim N^{\frac{p}{2}-1-\alpha(d)} \|P_N F(u)\|_{L_t^\infty L_x^2} + N^{p-\alpha(d)} \|P_N F(u)\|_{L_t^\infty L_x^{\frac{2d}{d+2+p}}} \\ &\lesssim N^{p-\alpha(d)} \|P_N F(u)\|_{L_t^\infty L_x^{\frac{2d}{d+2+p}}}. \end{aligned} \quad (6.6)$$

Using the Fundamental Theorem of Calculus, we decompose

$$\begin{aligned} F(u) &= O(|u_{>N_0}| |u_{\leq N_0}|^p) + O(|u_{>N_0}|^{1+p}) + F(u_{\frac{N}{10} \leq \cdot \leq N_0}) \\ &\quad + u_{< \frac{N}{10}} \int_0^1 F_z(u_{\frac{N}{10} \leq \cdot \leq N_0} + \theta u_{< \frac{N}{10}}) d\theta \\ &\quad + \overline{u_{< \frac{N}{10}}} \int_0^1 F_{\bar{z}}(u_{\frac{N}{10} \leq \cdot \leq N_0} + \theta u_{< \frac{N}{10}}) d\theta. \end{aligned} \quad (6.7)$$

The contribution to the right-hand side of (6.6) coming from terms that contain at least one copy of $u_{>N_0}$ can be estimated in the following manner: Using Hölder, Bernstein, Sobolev embedding, and (6.1),

$$\begin{aligned} N^{p-\alpha(d)} \|P_N O(|u_{>N_0}| |u|^p)\|_{L_t^\infty L_x^{\frac{2d}{d+2+p}}} &\lesssim N^{p-\alpha(d)} \|u\|_{L_t^\infty L_x^{\frac{dp}{2}}}^p \|u_{>N_0}\|_{L_t^\infty L_x^{\frac{2d}{d-2+p}}} \\ &\lesssim_u N^{p-\alpha(d)} N_0^{\alpha(d)-p}. \end{aligned} \quad (6.8)$$

Thus, this contribution is acceptable.

Next we turn to the contribution to the right-hand side of (6.6) coming from the last two terms in (6.7); it suffices to consider the first of them since similar arguments can be used to deal with the second.

First we note that as $|\nabla|^{s_c} u \in L_t^\infty L_x^2$, by Sobolev embedding we must have $\nabla u \in L_t^\infty L_x^{\frac{dp}{2+p}}$. Thus, an application of Lemma 2.9 together with (6.1) yield

$$\|P_{> \frac{N}{10}} F_z(u)\|_{L_t^\infty L_x^{\frac{d}{2+p}}} \lesssim N^{-p} \|\nabla u\|_{L_t^\infty L_x^{\frac{dp}{2+p}}}^p \lesssim N^{-p} \| |\nabla|^{s_c} u \|_{L_t^\infty L_x^2}^p \lesssim_u N^{-p}.$$

Thus, by Hölder's inequality and (6.3),

$$\begin{aligned} N^{p-\alpha(d)} &\left\| P_N \left(u_{< \frac{N}{10}} \int_0^1 F_z(u_{\frac{N}{10} \leq \cdot \leq N_0} + \theta u_{< \frac{N}{10}}) d\theta \right) \right\|_{L_t^\infty L_x^{\frac{2d}{d+2+p}}} \\ &\lesssim N^{p-\alpha(d)} \|u_{< \frac{N}{10}}\|_{L_t^\infty L_x^{\frac{2d}{d-2-p}}} \left\| P_{> \frac{N}{10}} \left(\int_0^1 F_z(u_{\frac{N}{10} \leq \cdot \leq N_0} + \theta u_{< \frac{N}{10}}) d\theta \right) \right\|_{L_t^\infty L_x^{\frac{d}{2+p}}} \end{aligned}$$

$$\begin{aligned}
&\lesssim N^{-\alpha(d)} \|u_{<\frac{N}{10}}\|_{L_t^\infty L_x^{\frac{2d}{d-2-p}}} \| |\nabla|^{s_c} u_{\leq N_0} \|_{L_t^\infty L_x^2}^p \\
&\lesssim_u \eta^p \sum_{N_1 < \frac{N}{10}} \left(\frac{N}{N_1}\right)^{\alpha(d)} A(N_1).
\end{aligned} \tag{6.9}$$

Hence, the contribution coming from the last two terms in (6.7) is acceptable.

We are left to estimate the contribution of $F(u_{\frac{N}{10} \leq \cdot \leq N_0})$ to the right-hand side of (6.6). We need only show

$$\|F(u_{\frac{N}{10} \leq \cdot \leq N_0})\|_{L_t^\infty L_x^{\frac{2d}{d+2+p}}} \lesssim_u \eta^p \sum_{\frac{N}{10} \leq N_1 \leq N_0} N_1^{\alpha(d)-p} A(N_1). \tag{6.10}$$

As $d \geq 7$ and the critical regularity s_c satisfies (1.5), we must have $p < 1$. Using the triangle inequality, Bernstein, (6.3), and Hölder, we estimate

$$\begin{aligned}
&\|F(u_{\frac{N}{10} \leq \cdot \leq N_0})\|_{L_t^\infty L_x^{\frac{2d}{d+2+p}}} \\
&\lesssim \sum_{\frac{N}{10} \leq N_1 \leq N_0} \|u_{N_1} |u_{\frac{N}{10} \leq \cdot \leq N_0}|^p\|_{L_t^\infty L_x^{\frac{2d}{d+2+p}}} \\
&\lesssim \sum_{\frac{N}{10} \leq N_1, N_2 \leq N_0} \|u_{N_1} |u_{N_2}|^p\|_{L_t^\infty L_x^{\frac{2d}{d+2+p}}} \\
&\lesssim \sum_{\frac{N}{10} \leq N_1 \leq N_2 \leq N_0} \|u_{N_1}\|_{L_t^\infty L_x^{\frac{2d}{d-2-p}}} \|u_{N_2}\|_{L_t^\infty L_x^{\frac{dp}{2+p}}}^p \\
&\quad + \sum_{\frac{N}{10} \leq N_2 \leq N_1 \leq N_0} \|u_{N_1}\|_{L_t^\infty L_x^{\frac{dp}{2+p}}}^p \|u_{N_1}\|_{L_t^\infty L_x^{\frac{2d}{d-2-p}}}^{1-p} \|u_{N_2}\|_{L_t^\infty L_x^{\frac{2d}{d-2-p}}}^p \\
&\lesssim \sum_{\frac{N}{10} \leq N_1 \leq N_2 \leq N_0} \|u_{N_1}\|_{L_t^\infty L_x^{\frac{2d}{d-2-p}}} N_2^{-p} \| |\nabla|^{s_c} u_{N_2} \|_{L_t^\infty L_x^2}^p \\
&\quad + \sum_{\frac{N}{10} \leq N_2 \leq N_1 \leq N_0} N_1^{-p} \| |\nabla|^{s_c} u_{N_1} \|_{L_t^\infty L_x^2}^p \|u_{N_1}\|_{L_t^\infty L_x^{\frac{2d}{d-2-p}}}^{1-p} \|u_{N_2}\|_{L_t^\infty L_x^{\frac{2d}{d-2-p}}}^p \\
&\lesssim_u \eta^p \sum_{\frac{N}{10} \leq N_1 \leq N_0} N_1^{\alpha(d)-p} A(N_1) \\
&\quad + \eta^p \sum_{\frac{N}{10} \leq N_2 \leq N_1 \leq N_0} \left(\frac{N_2}{N_1}\right)^{p^2} [N_1^{\alpha(d)-p} A(N_1)]^{1-p} [N_2^{\alpha(d)-p} A(N_2)]^p \\
&\lesssim_u \eta^p \sum_{\frac{N}{10} \leq N_1 \leq N_0} N_1^{\alpha(d)-p} A(N_1).
\end{aligned}$$

This proves (6.10) and so completes the proof of the lemma in dimensions $d \geq 7$.

Consider now $d = 5, 6$. Note that in this case, our assumptions guarantee that $1 < p < 2$. Arguing as for (6.6), we have

$$N^{-\alpha(d)} \|u_N(0)\|_{L_x^{\frac{2d}{d-3}}} \lesssim N^{1-\alpha(d)} \|P_N F(u)\|_{L_t^\infty L_x^{\frac{2d}{d+3}}},$$

which we estimate by decomposing the nonlinearity as in (6.7). The analogue of (6.8) in this case is

$$N^{1-\alpha(d)} \|P_N O(|u_{>N_0}| |u|^p)\|_{L_t^\infty L_x^{\frac{2d}{d+3}}} \lesssim N^{1-\alpha(d)} \|u\|_{L_t^\infty L_x^{\frac{dp}{2}}}^p \|u_{>N_0}\|_{L_t^\infty L_x^{\frac{2d}{d-1}}} \lesssim_u \left(\frac{N}{N_0}\right)^{1-\alpha(d)}.$$

Using Bernstein and Lemma 2.5 together with (6.3) and Sobolev embedding, we replace (6.9) by

$$\begin{aligned}
& N^{1-\alpha(d)} \left\| P_N \left(u_{<\frac{N}{10}} \int_0^1 F_z(u_{\frac{N}{10} \leq \cdot \leq N_0} + \theta u_{<\frac{N}{10}}) d\theta \right) \right\|_{L_t^\infty L_x^{\frac{2d}{d+3}}} \\
& \lesssim N^{1-\alpha(d)} \|u_{<\frac{N}{10}}\|_{L_t^\infty L_x^{\frac{2d}{d+3}}} \|P_{>\frac{N}{10}} \left(\int_0^1 F_z(u_{\frac{N}{10} \leq \cdot \leq N_0} + \theta u_{<\frac{N}{10}}) d\theta \right)\|_{L_t^\infty L_x^{\frac{d}{3}}} \\
& \lesssim N^{-\alpha(d)} \|u_{<\frac{N}{10}}\|_{L_t^\infty L_x^{\frac{2d}{d+3}}} \|\nabla u_{\leq N_0}\|_{L_t^\infty L_x^{\frac{dp}{2+p}}} \|u_{\leq N_0}\|_{L_t^\infty L_x^{\frac{dp}{2}}}^{p-1} \\
& \lesssim \sum_{N_1 < \frac{N}{10}} \left(\frac{N_1}{N}\right)^{\alpha(d)} A(N_1) \|\nabla|^{s_c} u_{\leq N_0}\|_{L_t^\infty L_x^2}^p \\
& \lesssim_u \eta^p \sum_{N_1 < \frac{N}{10}} \left(\frac{N_1}{N}\right)^{\alpha(d)} A(N_1).
\end{aligned}$$

Finally, arguing as for (6.10), we estimate

$$\begin{aligned}
& \|F(u_{\frac{N}{10} \leq \cdot \leq N_0})\|_{L_t^\infty L_x^{\frac{2d}{d+3}}} \\
& \lesssim \sum_{\frac{N}{10} \leq N_1, N_2 \leq N_0} \|u_{N_1} u_{N_2} |u_{\frac{N}{10} \leq \cdot \leq N_0}|^{p-1}\|_{L_t^\infty L_x^{\frac{2d}{d+3}}} \\
& \lesssim \sum_{\frac{N}{10} \leq N_1 \leq N_2, N_3 \leq N_0} \|u_{N_1}\|_{L_t^\infty L_x^{\frac{2d}{d+3}}} \|u_{N_2}\|_{L_t^\infty L_x^{\frac{dp}{3}}} \|u_{N_3}\|_{L_t^\infty L_x^{\frac{dp}{3}}}^{p-1} \\
& \quad + \sum_{\frac{N}{10} \leq N_3 \leq N_1 \leq N_2 \leq N_0} \|u_{N_1}\|_{L_t^\infty L_x^{\frac{2d}{d+3}}}^{2-p} \|u_{N_1}\|_{L_t^\infty L_x^{\frac{dp}{3}}}^{p-1} \|u_{N_2}\|_{L_t^\infty L_x^{\frac{dp}{3}}} \|u_{N_3}\|_{L_t^\infty L_x^{\frac{dp}{3}}}^{p-1} \\
& \lesssim_u \sum_{\frac{N}{10} \leq N_1 \leq N_2, N_3 \leq N_0} \|u_{N_1}\|_{L_t^\infty L_x^{\frac{2d}{d+3}}} \eta N_2^{-\frac{1}{p}} (\eta N_3^{-\frac{1}{p}})^{p-1} \\
& \quad + \sum_{\frac{N}{10} \leq N_3 \leq N_1 \leq N_2 \leq N_0} \|u_{N_1}\|_{L_t^\infty L_x^{\frac{2d}{d+3}}}^{2-p} (\eta N_1^{-\frac{1}{p}})^{p-1} \eta N_2^{-\frac{1}{p}} \|u_{N_3}\|_{L_t^\infty L_x^{\frac{dp}{3}}}^{p-1} \\
& \lesssim_u \eta^p \sum_{\frac{N}{10} \leq N_1 \leq N_0} N_1^{\alpha(d)-1} A(N_1) \\
& \quad + \eta^p \sum_{\frac{N}{10} \leq N_3 \leq N_1 \leq N_0} \left(\frac{N_3}{N_1}\right)^{p-1} [N_1^{\alpha(d)-1} A(N_1)]^{2-p} [N_3^{\alpha(d)-1} A(N_3)]^{p-1} \\
& \lesssim_u \eta^p \sum_{\frac{N}{10} \leq N_1 \leq N_0} N_1^{\alpha(d)-1} A(N_1).
\end{aligned}$$

Putting everything together completes the proof of the lemma when $d = 5, 6$. $\square$

This lemma leads very quickly to our first goal:

Proposition 6.3 (L^p breach of scaling). *Let u be as in Theorem 6.1. Then*

$$u \in L_t^\infty L_x^q \quad \text{for some } 2 < q < \frac{dp}{2}. \quad (6.11)$$

In particular, for $0 \leq s \leq s_c$,

$$|\nabla|^s u \in L_t^\infty L_x^2 \implies |\nabla|^s F(u) \in L_t^\infty L_x^r \quad \text{for some } r < \frac{2d}{d+4}. \quad (6.12)$$

Proof. We first consider (6.11). We will only present the details for $d \geq 7$. The treatment of $d = 5, 6$ is completely analogous.

Combining Lemma 6.2 with Lemma 2.13, we deduce

$$\|u_N\|_{L_t^\infty L_x^{\frac{2d}{d-2-p}}} \lesssim_u N^{p-} \quad \text{for all } N \leq 10N_0. \quad (6.13)$$

In applying Lemma 2.13, we set $N = 10 \cdot 2^{-k} N_0$, $x_k = A(10 \cdot 2^{-k} N_0)$, and take η sufficiently small. Note that $x_k \in \ell^\infty$ by virtue of (6.4).

By interpolation followed by (6.13), Bernstein, and (6.1),

$$\|u_N\|_{L_t^\infty L_x^q} \lesssim \|u_N\|_{L_t^\infty L_x^{\frac{2d}{d-2-p}}}^{\frac{d(q-2)}{q(2+p)}} \|u_N\|_{L_t^\infty L_x^2}^{\frac{2d-q(d-2-p)}{q(2+p)}} \lesssim_u N^{\frac{dp(q-2)}{q(2+p)}} N^{-\left(\frac{d}{2}-\frac{2}{p}\right)\frac{2d-q(d-2-p)}{q(2+p)}}$$

for all $N \leq 10N_0$. Note that the power of N appearing in the formula above is positive for

$$\frac{2d(2p^2+dp-4)}{dp^2+(d^2-2d+4)p-4(d-2)} < q < \frac{dp}{2}.$$

Thus, for these values of q , Bernstein together with (6.1) yield

$$\|u\|_{L_t^\infty L_x^q} \leq \|u_{\leq N_0}\|_{L_t^\infty L_x^q} + \|u_{> N_0}\|_{L_t^\infty L_x^q} \lesssim_u \sum_{N \leq N_0} N^{0+} + \sum_{N > N_0} N^{\frac{2}{p}-\frac{d}{q}} \lesssim_u 1,$$

which completes the proof of (6.11).

We turn now to (6.12). For $0 \leq s \leq 1$, we use Lemma 2.5 to estimate

$$\| |\nabla|^s F(u) \|_{L_t^\infty L_x^r} \lesssim \| |\nabla|^s u \|_{L_t^\infty L_x^2} \|u\|_{L_t^\infty L_x^{\frac{2rp}{2-r}}}^p.$$

We will return to this shortly, but first we consider the case $1 < s \leq s_c$. Using Lemma 2.4 together with Sobolev embedding,

$$\begin{aligned} \| |\nabla|^s F(u) \|_{L_t^\infty L_x^r} &\lesssim \| |\nabla|^{s-1} (\nabla u \cdot F'(u)) \|_{L_t^\infty L_x^r} \\ &\lesssim \| |\nabla|^s u \|_{L_t^\infty L_x^2} \|u\|_{L_t^\infty L_x^{\frac{2rp}{2-r}}}^p + \|\nabla u\|_{L_t^\infty L_x^{\frac{dp}{2+p}}} \| |\nabla|^{s-1} F'(u) \|_{L_t^\infty L_x^{\frac{dpr}{dp-r(2+p)}}} \\ &\lesssim \| |\nabla|^s u \|_{L_t^\infty L_x^2} \left(\|u\|_{L_t^\infty L_x^{\frac{2rp}{2-r}}}^p + \| |\nabla|^{s-1} F'(u) \|_{L_t^\infty L_x^{\frac{dpr}{dp-r(2+p)}}} \right). \end{aligned}$$

For $p \geq 1$, we invoke Lemma 2.5 and then use Sobolev embedding to estimate

$$\begin{aligned} \| |\nabla|^{s-1} F'(u) \|_{L_t^\infty L_x^{\frac{dpr}{dp-r(2+p)}}} &\lesssim \| |\nabla|^{s-1} u \|_{L_t^\infty L_x^{\frac{2d}{d-2}}} \|u\|_{L_t^\infty L_x^{\frac{2dpr(p-1)}{2dp-dpr-4r}}}^{p-1} \\ &\lesssim \| |\nabla|^s u \|_{L_t^\infty L_x^2} \|u\|_{L_t^\infty L_x^{\frac{2dpr(p-1)}{2dp-dpr-4r}}}^{p-1}. \end{aligned}$$

If instead $p < 1$, we use Lemma 2.6 and Sobolev embedding to obtain

$$\begin{aligned} \| |\nabla|^{s-1} F'(u) \|_{L_t^\infty L_x^{\frac{dpr}{dp-r(2+p)}}} &\lesssim \| |\nabla|^\sigma u \|_{L_t^\infty L_x^{\frac{dp}{2+\sigma p}}}^{\frac{s-1}{\sigma}} \|u\|_{L_t^\infty L_x^{\frac{dpr(\sigma p+1-s)}{dp\sigma-\sigma prs-2r(s-1)-2\sigma r}}}^{p-\frac{s-1}{\sigma}} \\ &\lesssim \| |\nabla|^s u \|_{L_t^\infty L_x^2}^{\frac{s-1}{\sigma}} \|u\|_{L_t^\infty L_x^{\frac{dpr(\sigma p+1-s)}{dp\sigma-\sigma prs-2r(s-1)-2\sigma r}}}^{p-\frac{s-1}{\sigma}} \end{aligned}$$

for some $\frac{s-1}{p} < \sigma < 1$.

For all $0 \leq s \leq s_c$, to obtain (6.12) it thus suffices to note that for $r < \frac{2d}{d+4}$ the exponents $\frac{2rp}{2-r}$, $\frac{2dpr(p-1)}{2dp-dpr-4r}$, $\frac{dpr(\sigma p+1-s)}{dp\sigma-\sigma prs-2r(s-1)-2\sigma r}$ are all less than $\frac{dp}{2}$ and converge to this number as $r \rightarrow \frac{2d}{d+4}$. Thus, (6.11) yields the claim. $\square$

Following [19], our second step is to use the double Duhamel trick to upgrade (6.11) to a negative regularity statement in L_x^2 -based Sobolev spaces. The arguments we present follow closely the ones in [19]. For the sake of completeness, we present the details below.

Proposition 6.4 (Some negative regularity). *Let $d \geq 5$ and assume the critical regularity s_c obeys (1.5). Let u be as in Theorem 6.1. Assume further that $|\nabla|^s F(u) \in L_t^\infty L_x^r$ for some $r < \frac{2d}{d+4}$ and some $0 \leq s \leq s_c$. Then there exists $s_0 = s_0(d, r) > 0$ such that $u \in L_t^\infty \dot{H}_x^{s-s_0+}$.*

Proof. The proposition will follow once we establish

$$\| |\nabla|^s u_N \|_{L_t^\infty L_x^2} \lesssim_u N^{s_0} \quad \text{for all } N > 0 \quad \text{and} \quad s_0 := \frac{d}{r} - \frac{d+4}{2} > 0. \quad (6.14)$$

Indeed, by Bernstein combined with (6.1),

$$\begin{aligned} \| |\nabla|^{s-s_0+} u \|_{L_t^\infty L_x^2} &\leq \| |\nabla|^{s-s_0+} u_{\leq 1} \|_{L_t^\infty L_x^2} + \| |\nabla|^{s-s_0+} u_{>1} \|_{L_t^\infty L_x^2} \\ &\lesssim_u \sum_{N \leq 1} N^{0+} + \sum_{N > 1} N^{(s-s_0+)-s_c} \\ &\lesssim_u 1. \end{aligned}$$

Thus, we are left to prove (6.14). By time-translation symmetry, it suffices to prove

$$\| |\nabla|^s u_N(0) \|_{L_x^2} \lesssim_u N^{s_0} \quad \text{for all } N > 0 \quad \text{and} \quad s_0 := \frac{d}{r} - \frac{d+4}{2} > 0. \quad (6.15)$$

Using the Duhamel formula (1.7) both in the future and in the past, we write

$$\begin{aligned} &\| |\nabla|^s u_N(0) \|_{L_x^2}^2 \\ &= \lim_{T \rightarrow \infty} \lim_{T' \rightarrow -\infty} \left\langle i \int_0^T e^{-it\Delta} P_N |\nabla|^s F(u(t)) dt, -i \int_{T'}^0 e^{-i\tau\Delta} P_N |\nabla|^s F(u(\tau)) d\tau \right\rangle \\ &\leq \int_0^\infty \int_{-\infty}^0 \left| \langle P_N |\nabla|^s F(u(t)), e^{i(t-\tau)\Delta} P_N |\nabla|^s F(u(\tau)) \rangle \right| dt d\tau. \end{aligned}$$

We estimate the term inside the integrals in two ways. On one hand, using Hölder and the dispersive estimate,

$$\begin{aligned} &\left| \langle P_N |\nabla|^s F(u(t)), e^{i(t-\tau)\Delta} P_N |\nabla|^s F(u(\tau)) \rangle \right| \\ &\lesssim \| P_N |\nabla|^s F(u(t)) \|_{L_x^r} \| e^{i(t-\tau)\Delta} P_N |\nabla|^s F(u(\tau)) \|_{L_{x'}^{r'}} \\ &\lesssim |t - \tau|^{\frac{d}{2} - \frac{d}{r}} \| |\nabla|^s F(u) \|_{L_t^\infty L_x^r}^2. \end{aligned}$$

On the other hand, using Bernstein,

$$\begin{aligned} &\left| \langle P_N |\nabla|^s F(u(t)), e^{i(t-\tau)\Delta} P_N |\nabla|^s F(u(\tau)) \rangle \right| \\ &\lesssim \| P_N |\nabla|^s F(u(t)) \|_{L_x^2} \| e^{i(t-\tau)\Delta} P_N |\nabla|^s F(u(\tau)) \|_{L_x^2} \\ &\lesssim N^{2(\frac{d}{2} - \frac{d}{r})} \| |\nabla|^s F(u) \|_{L_t^\infty L_x^r}^2. \end{aligned}$$

Thus,

$$\begin{aligned} \| |\nabla|^s u_N(0) \|_{L_x^2}^2 &\lesssim \| |\nabla|^s F(u) \|_{L_t^\infty L_x^r}^2 \int_0^\infty \int_{-\infty}^0 \min\{|t - \tau|^{-1}, N^2\}^{\frac{d}{r} - \frac{d}{2}} dt d\tau \\ &\lesssim N^{2s_0} \| |\nabla|^s F(u) \|_{L_t^\infty L_x^r}^2. \end{aligned}$$

To obtain the last inequality we used the fact that $\frac{d}{r} - \frac{d}{2} > 2$ since $r < \frac{2d}{d+4}$. Thus (6.15) holds; this finishes the proof of the proposition. $\square$

The proof of Theorem 6.1 will follow from iterating Proposition 6.4 finitely many times.

Proof of Theorem 6.1. Proposition 6.3 allows us to apply Proposition 6.4 with $s = s_c$. We conclude that $u \in L_t^\infty \dot{H}_x^{s_c-s_0+}$ for some $s_0 = s_0(d, r) > 0$. Combining this with (6.12), we deduce that $|\nabla|^{s_c-s_0+} F(u) \in L_t^\infty L_x^r$ for some $r < \frac{2d}{d+4}$. We are thus in the position to apply

Proposition 6.4 again and obtain $u \in L_t^\infty \dot{H}_x^{s_c-2s_0+}$. Iterating this procedure finitely many times, we derive $u \in L_t^\infty \dot{H}_x^{-\varepsilon}$ for any $0 < \varepsilon < s_0$.

This completes the proof of Theorem 6.1. $\square$

To prove the equivalent of Theorem 6.1 in the setting of Theorem 1.4, one repeats the arguments presented above with

$$A(N) := N^{\frac{d-5}{2}} \sup_{t \in \mathbb{R}} \|u_N(t)\|_{L_x^{\frac{2d}{d-3}}}.$$

7. THE LOW-TO-HIGH FREQUENCY CASCADE

In this section, we use the negative regularity proved in the previous section to preclude low-to-high frequency cascade solutions.

Theorem 7.1 (Absence of cascades). *In the settings of Theorems 1.4 and 1.5 there are no solutions to (1.1) that are low-to-high frequency cascades in the sense of Theorem 1.13.*

Proof. Suppose for a contradiction that there existed such a solution u . Then, by the results proved in Section 6, $u \in L_t^\infty L_x^2$; thus, by the conservation of mass,

$$0 \leq M(u) = M(u(t)) = \int_{\mathbb{R}^d} |u(t, x)|^2 dx < \infty \quad \text{for all } t \in \mathbb{R}.$$

Fix $t \in \mathbb{R}$ and let $\eta > 0$ be a small constant. By compactness (see Remark 1.11),

$$\int_{|\xi| \leq c(\eta)N(t)} |\xi|^{2s_c} |\hat{u}(t, \xi)|^2 d\xi \leq \eta.$$

On the other hand, as $u \in L_t^\infty \dot{H}_x^{-\varepsilon}$ for some $\varepsilon > 0$,

$$\int_{|\xi| \leq c(\eta)N(t)} |\xi|^{-2\varepsilon} |\hat{u}(t, \xi)|^2 d\xi \lesssim_u 1.$$

Hence, by Hölder's inequality,

$$\int_{|\xi| \leq c(\eta)N(t)} |\hat{u}(t, \xi)|^2 d\xi \lesssim_u \eta^{\frac{\varepsilon}{s_c + \varepsilon}}. \quad (7.1)$$

Meanwhile, by elementary considerations and $u \in L_t^\infty \dot{H}_x^{s_c}$,

$$\begin{aligned} \int_{|\xi| \geq c(\eta)N(t)} |\hat{u}(t, \xi)|^2 d\xi &\leq [c(\eta)N(t)]^{-2s_c} \int_{\mathbb{R}^d} |\xi|^{2s_c} |\hat{u}(t, \xi)|^2 d\xi \\ &\leq [c(\eta)N(t)]^{-2s_c} \|\nabla|^{s_c} u(t)\|_2^2 \\ &\lesssim_u [c(\eta)N(t)]^{-2s_c}. \end{aligned} \quad (7.2)$$

Collecting (7.1) and (7.2) and using Plancherel's theorem, we obtain

$$0 \leq M(u) \lesssim_u c(\eta)^{-2s_c} N(t)^{-2s_c} + \eta^{\frac{\varepsilon}{s_c + \varepsilon}}$$

for all $t \in \mathbb{R}$. As u is a low-to-high cascade, there is a sequence of times $t_n \rightarrow \infty$ so that $N(t_n) \rightarrow \infty$. As $\eta > 0$ is arbitrary, we may conclude $M(u) = 0$ and hence u is identically zero. This contradicts the fact that $S_{\mathbb{R}}(u) = \infty$, thus settling Theorem 7.1. $\square$

8. THE SOLITON

In this section, we use the negative regularity proved in Section 6 to preclude soliton-like solutions.

Theorem 8.1 (Absence of solitons). *In the settings of Theorems 1.4 and 1.5 there are no global solutions to (1.1) that are solitons in the sense of Theorem 1.13.*

Proof. As usual, we will use a monotonicity formula to preclude soliton-like solutions. Since we are in the defocusing case, we will use the interaction Morawetz inequality introduced in [7]. For the high dimensional case discussed here, the details of this derivation can be found in [33] or [37].

To prove Theorem 8.1, we argue by contradiction. We assume there exists a solution u to (1.1) which is a soliton in the sense of Theorem 1.13. Then, by the negative regularity results proved in Section 6, $u \in L_t^\infty H_x^1$. The interaction Morawetz inequality yields

$$\int_I \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{|u(t, x)|^2 |u(t, y)|^2}{|x - y|^3} dx dy dt \lesssim \|u\|_{L_t^\infty L_x^2(I \times \mathbb{R}^d)}^3 \|\nabla u\|_{L_t^\infty L_x^2(I \times \mathbb{R}^d)} \lesssim_u 1$$

for any compact time interval $I \subset \mathbb{R}$. As in dimension d convolution with $|x|^{-3}$ is basically the same as the fractional integration operator $|\nabla|^{-(d-3)}$, the interaction Morawetz inequality yields

$$\| |\nabla|^{-\frac{d-3}{2}} (|u|^2) \|_{L_{t,x}^2(I \times \mathbb{R}^d)} \lesssim_u 1.$$

By [37, Lemma 5.6], this implies

$$\| |\nabla|^{-\frac{d-3}{4}} u \|_{L_{t,x}^4(I \times \mathbb{R}^d)} \lesssim_u 1.$$

Interpolating between this estimate and the fact that $u \in L_t^\infty \dot{H}_x^1$, we derive

$$\|u\|_{L_t^{d+1} L_x^{\frac{2(d+1)}{d-1}}(I \times \mathbb{R}^d)} \lesssim_u 1 \quad (8.1)$$

for any compact time interval $I \subset \mathbb{R}$.

Next we claim that

$$\|u(t)\|_{L_x^{\frac{2(d+1)}{d-1}}} \gtrsim_u 1 \quad \text{uniformly for } t \in \mathbb{R}. \quad (8.2)$$

Otherwise, there exists a time sequence t_n such that $u(t_n)$ converges weakly to zero in $L_x^{\frac{2(d+1)}{d-1}}$. As $u(t)$ is uniformly bounded in $\dot{H}_x^{sc}$, this implies that $u(t_n)$ converges weakly to zero in $\dot{H}_x^{sc}$. As the orbit of u is precompact in $\dot{H}_x^{sc}$ and u is not identically zero, we derive a contradiction.

Using (8.1) and (8.2), we easily derive a contradiction by taking the interval I to be sufficiently long. $\square$

REFERENCES

- [1] P. Begout and A. Vargas, *Mass concentration phenomena for the L^2 -critical nonlinear Schrödinger equation*. Trans. Amer. Math. Soc. **359** (2007), 5257–5282. MR2327030
- [2] J. Bourgain, *Global wellposedness of defocusing critical nonlinear Schrödinger equation in the radial case*. J. Amer. Math. Soc. **12** (1999), 145–171. MR1626257
- [3] R. Carles and S. Keraani, *On the role of quadratic oscillations in nonlinear Schrödinger equation II. The L^2 -critical case*. Trans. Amer. Math. Soc. **359** (2007), 33–62.
- [4] T. Cazenave and F. B. Weissler, *The Cauchy problem for the critical nonlinear Schrödinger equation in H^s* . Nonlinear Anal. **14** (1990), 807–836. MR1055532
- [5] T. Cazenave, *Semilinear Schrödinger equations*. Courant Lecture Notes in Mathematics, **10**. American Mathematical Society, 2003. MR2002047
- [6] M. Christ and M. Weinstein, *Dispersion of small amplitude solutions of the generalized Korteweg-de Vries equation*. J. Funct. Anal. **100** (1991), 87–109. MR1124294

- [7] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao, *Global existence and scattering for rough solutions of a nonlinear Schrödinger equation on $\mathbb{R}^3$* . Comm. Pure Appl. Math. **57** (2004), 987–1014. MR2053757
- [8] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao, *Global well-posedness and scattering for the energy-critical nonlinear Schrödinger equation in $\mathbb{R}^3$* . Ann. of Math. **167** (2008), 767–865. MR2415387
- [9] L. Escauriaza, G. A. Seregin, V. Sverak, *$L_{3,\infty}$ -solutions of Navier-Stokes equations and backward uniqueness*. Uspekhi Mat. Nauk **58** (2003), 3–44; translation in Russian Math. Surveys **58** (2003), 211–250. MR1992563
- [10] J. Ginibre and G. Velo, *Smoothing properties and retarded estimates for some dispersive evolution equations*. Comm. Math. Phys. **144** (1992), 163–188. MR1151250
- [11] G. Grillakis, *On nonlinear Schrödinger equations*. Comm. PDE **25** (2000), 1827–1844. MR1778782
- [12] M. Keel and T. Tao, *Endpoint Strichartz Estimates*. Amer. J. Math. **120** (1998), 955–980. MR1646048
- [13] C. E. Kenig and F. Merle, *Global well-posedness, scattering and blow up for the energy-critical, focusing, nonlinear Schrödinger equation in the radial case*. Invent. Math. **166** (2006), 645–675. MR2257393
- [14] C. E. Kenig and F. Merle, *Scattering for $H^{1/2}$ bounded solutions to the cubic, defocusing NLS in 3 dimensions*. Preprint [arXiv:0712.1834](#).
- [15] C. E. Kenig and F. Merle, *Nondispersive radial solutions to energy supercritical non-linear wave equations, with applications*. Preprint [arXiv:0810.4834](#).
- [16] S. Keraani, *On the defect of compactness for the Strichartz estimates for the Schrödinger equations*. J. Diff. Eq. **175** (2001), 353–392. MR1855973
- [17] S. Keraani, *On the blow-up phenomenon of the critical nonlinear Schrödinger equation*. J. Funct. Anal. **235** (2006), 171–192. MR2216444
- [18] R. Killip, T. Tao, and M. Visan, *The cubic nonlinear Schrödinger equation in two dimensions with radial data*. To appear in JEMS. Preprint [arXiv:0707.3188](#).
- [19] R. Killip and M. Visan, *The focusing energy-critical nonlinear Schrödinger equation in dimensions five and higher*. Preprint [arXiv:0804.1018](#).
- [20] R. Killip and M. Visan, *Nonlinear Schrödinger equations at critical regularity*. Lecture notes prepared for Clay Mathematics Institute Summer School, Zürich, Switzerland, 2008.
- [21] R. Killip, M. Visan, and X. Zhang, *The mass-critical nonlinear Schrödinger equation with radial data in dimensions three and higher*. To appear in Analysis and PDE. Preprint [arXiv:0708.0849](#).
- [22] F. Merle and L. Vega, *Compactness at blow-up time for L^2 solutions of the critical nonlinear Schrödinger equation in 2D*. Internat. Math. Res. Notices **8** (1998), 399–425. MR1628235
- [23] K. Nakanishi, *Scattering theory for nonlinear Klein-Gordon equation with Sobolev critical power*. Internat. Math. Res. Notices **1** (1999), 31–60. MR1666973
- [24] E. Ryckman and M. Visan, *Global well-posedness and scattering for the defocusing energy-critical nonlinear Schrödinger equation in $\mathbb{R}^{1+4}$* . Amer. J. Math. **129** (2007), 1–60. MR2288737
- [25] S. Shao, *Maximizers for the Strichartz inequalities and the Sobolev–Strichartz inequalities for the Schrödinger equation*. Preprint [arXiv:0809.0153](#).
- [26] R. S. Strichartz, *Multipliers on fractional Sobolev spaces*. J. Math. Mech. **16** (1967), 1031–1060. MR0215084
- [27] R. S. Strichartz, *Restriction of Fourier transform to quadratic surfaces and decay of solutions of wave equations*. Duke Math. J. **44** (1977), 705–774. MR0512086
- [28] T. Tao, *Global well-posedness and scattering for the higher-dimensional energy-critical non-linear Schrödinger equation for radial data*. New York J. of Math. **11** (2005), 57–80. MR2154347
- [29] T. Tao, *A (concentration-)compact attractor for high-dimensional non-linear Schrödinger equations*. Dyn. Partial Differ. Equ. **4** (2007), 1–53. MR2304091
- [30] T. Tao, *Nonlinear dispersive equations. Local and global analysis*. CBMS Regional Conference Series in Mathematics, 106. American Mathematical Society, Providence, RI, 2006. MR2233925
- [31] T. Tao, *Why global regularity for Navier–Stokes is hard*. In “Structure and randomness: pages from year one of a mathematical blog”. American Mathematical Society, 2008.
- [32] T. Tao and M. Visan, *Stability of energy-critical nonlinear Schrödinger equations in high dimensions*. Electron. J. Diff. Eqns. **118** (2005), 1–28. MR2174550
- [33] T. Tao, M. Visan, and X. Zhang, *The nonlinear Schrödinger equation with combined power-type nonlinearities*. Comm. Partial Differential Equations **32** (2007), 1281–1343. MR2354495
- [34] T. Tao, M. Visan, and X. Zhang, *Minimal-mass blowup solutions of the mass-critical NLS*. Forum Math. **20** (2008), 881–919. MR2445122
- [35] T. Tao, M. Visan, and X. Zhang, *Global well-posedness and scattering for the mass-critical nonlinear Schrödinger equation for radial data in high dimensions*. Duke Math. J. **140** (2007), 165–202. MR2355070
- [36] M. E. Taylor, *Tools for PDE*. Mathematical Surveys and Monographs, **81**. American Mathematical Society, Providence, RI, 2000. MR1766415

- [37] M. Visan, *The defocusing energy-critical nonlinear Schrödinger equation in higher dimensions*. Duke Math. J. **138** (2007), 281–374. MR2318286
- [38] M. Visan, *The defocusing energy-critical nonlinear Schrödinger equation in dimensions five and higher*. Ph.D. Thesis, UCLA, 2006.

UNIVERSITY OF CALIFORNIA, LOS ANGELES, CA

UNIVERSITY OF CHICAGO, CHICAGO, IL